


1st Congressional District Nutrient and Sediment Load Reductions


Accomplished By Private Landowners and the Indiana Conservation Partnership



Comparison of Landuse Across District




Land use calculated using the 2019 NASS Cropland Data Layer



- County Seats
- County Boundaries
- Congressional District

Sediment Reduced: 30,998,700 lbs.

Enough to fill 155 freight cars!



Phosphorus Reduced: 19,610 lbs.

Enough to fill 20 truck beds (8' bed)!



Nitrogen Reduced: 39,175 lbs.

Enough to fill 39 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2014	169	205	17,724,605	11,090	22,175
2015	153	219	16,467,295	10,085	20,180
2016	110	186	12,491,940	7,265	14,490
2017	131	233	18,786,670	11,680	23,310
2018	155	277	18,714,900	11,640	23,220
2019	319	491	30,998,700	19,610	39,175
13-18	1,164		115,184,110	79,620	159,035




The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 metrics are excluded from the table due to space limitations, but are present in the "13-18" summations.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact ISDANutrientReduction@isda.in.gov
Last updated: 3/16/2020


Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

2nd Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Sediment Reduced: 213,296,100 lbs.
 Enough to fill 1,066 freight cars!



Phosphorus Reduced: 114,795 lbs.
 Enough to fill 115 truck beds (8' bed)!



Nitrogen Reduced: 229,635 lbs.
 Enough to fill 230 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2014	2,008	2,274	268,418,595	150,565	301,520
2015	1,708	2,081	216,213,330	123,625	248,270
2016	1,435	1,969	189,528,870	106,680	213,670
2017	1,581	2,342	199,865,675	112,505	225,375
2018	1,366	2,267	169,637,610	96,070	192,410
2019	1,480	2,650	213,296,100	114,795	229,635
13-18	11,330		1,256,960,180	812,020	1,626,710



The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 metrics are excluded from the table due to space limitations, but are present in the "13-18" summations.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact ISDANutrientReduction@isda.in.gov
 Last updated: 3/16/2020


Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

3rd Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership




Comparison of Landuse Across District



Land use calculated using the 2019 NASS Cropland Data Layer

Sediment Reduced: 449,031,030 lbs.

Enough to fill 2,245 freight cars!



Phosphorus Reduced: 285,170 lbs.

Enough to fill 285 truck beds (8' bed)!



Nitrogen Reduced: 601,615 lbs.

Enough to fill 602 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2014	2,018	2,365	254,369,525	154,220	322,525
2015	2,448	3,031	309,059,770	190,600	395,075
2016	2,055	2,976	288,755,885	181,145	376,160
2017	2,526	3,596	324,149,870	201,450	416,745
2018	2,712	4,113	291,676,395	192,985	417,390
2019	3,615	5,384	449,031,030	285,170	601,615
13-18	18,366		1,917,042,475	1,400,260	2,932,690




The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 metrics are excluded from the table due to space limitations, but are present in the "13-18" summations.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact ISDANutrientReduction@isda.in.gov
 Last updated: 3/16/2020

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.


4th Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Sediment Reduced: 403,202,235 lbs.

Enough to fill 2,016 freight cars!



Phosphorus Reduced: 229,265 lbs.

Enough to fill 229 truck beds (8' bed)!



Nitrogen Reduced: 465,700 lbs.

Enough to fill 466 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2014	1,558	2,288	244,080,585	143,360	289,115
2015	1,751	3,020	295,670,170	171,210	344,910
2016	1,533	3,387	276,199,165	159,340	327,075
2017	1,727	4,073	335,472,520	193,605	394,955
2018	2,002	4,616	376,341,565	215,580	438,490
2019	1,870	5,020	403,202,235	229,265	465,700
13-18	12,232		1,930,966,240	1,215,530	2,467,345



The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 metrics are excluded from the table due to space limitations, but are present in the "13-18" summations.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact ISDANutrientReduction@isda.in.gov
 Last updated: 3/16/2020


Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

5th Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Sediment Reduced: 35,259,820 lbs.
 Enough to fill 176 freight cars!



Phosphorus Reduced: 22,750 lbs.
 Enough to fill 23 truck beds (8' bed)!



Nitrogen Reduced: 46,515 lbs.
 Enough to fill 47 truck beds (8' bed)!






Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2014	221	244	23,636,670	14,985	29,950
2015	189	268	27,069,660	18,160	37,350
2016	73	231	15,367,800	9,815	20,670
2017	142	321	23,487,525	14,795	30,615
2018	162	384	24,183,300	15,345	31,725
2019	206	476	35,259,820	22,750	46,515
13-18	1,189		149,004,775	110,075	225,245


The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 metrics are excluded from the table due to space limitations, but are present in the "13-18" summations.

6th Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Sediment Reduced: 491,485,490 lbs.
 Enough to fill 2,457 freight cars!



Phosphorus Reduced: 254,045 lbs.
 Enough to fill 254 truck beds (8' bed)!



Nitrogen Reduced: 508,875 lbs.
 Enough to fill 509 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2014	1,228	1,820	244,428,150	123,630	247,640
2015	1,511	2,338	295,058,725	148,135	296,720
2016	1,554	2,767	298,013,170	149,700	299,705
2017	1,476	3,122	302,656,570	154,055	308,410
2018	2,026	3,968	372,314,755	191,905	384,745
2019	2,656	5,233	491,485,490	254,045	508,875
13-18	12,182		2,003,956,860	1,137,185	2,277,635

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 metrics are excluded from the table due to space limitations, but are present in the "13-18" summations.

7th Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Boone

Hamilton

Hendricks

Marion

Hancock

Indianapolis

Morgan

Johnson


Shelby

● County Seats

□ County Boundaries

■ Congressional District

Comparison of Landuse Across District



Land use calculated using the 2019 NASS Cropland Data Layer

Sediment Reduced: 1,839,640 lbs.

Enough to fill 9 freight cars!



Phosphorus Reduced: 795 lbs.

Enough to fill 1 truck beds (8' bed)!



Nitrogen Reduced: 1,525 lbs.

Enough to fill 2 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance.

Nutrient estimates only consider sediment bound N and P, not dissolved components.

Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2014	17	22	2,050,150	940	1,815
2015	20	31	2,189,850	1,030	1,995
2016	33	45	2,018,970	920	1,770
2017	36	58	2,176,835	1,040	2,015
2018	29	57	1,717,210	725	1,380
2019	23	60	1,839,640	795	1,525
13-18	174		11,992,655	5,755	11,110



The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 metrics are excluded from the table due to space limitations, but are present in the "13-18" summations.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact ISDANutrientReduction@isda.in.gov
Last updated: 3/16/2020


Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

8th Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Sediment Reduced: 987,849,895 lbs.
 Enough to fill 4,939 freight cars!



Phosphorus Reduced: 470,400 lbs.
 Enough to fill 470 truck beds (8' bed)!



Nitrogen Reduced: 941,675 lbs.
 Enough to fill 942 truck beds (8' bed)!






Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2014	2,603	3,465	742,172,930	345,865	692,265
2015	2,730	4,047	856,394,375	400,300	801,075
2016	2,254	4,029	751,253,275	352,535	705,780
2017	2,426	4,739	834,775,050	391,060	783,085
2018	3,638	6,268	905,871,620	425,010	850,965
2019	3,308	7,105	987,849,895	470,400	941,675
13-18	20,088		5,078,317,145	2,728,500	5,461,850


The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 metrics are excluded from the table due to space limitations, but are present in the "13-18" summations.

9th Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Sediment Reduced: 716,146,085 lbs.
Enough to fill 3,581 freight cars!



Phosphorus Reduced: 336,410 lbs.
Enough to fill 336 truck beds (8' bed)!



Nitrogen Reduced: 675,745 lbs.
Enough to fill 676 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2014	1,340	1,835	410,683,580	189,375	378,840
2015	1,500	2,295	534,566,110	250,165	502,825
2016	1,650	2,850	569,237,190	264,675	531,875
2017	1,865	3,472	704,330,910	327,110	656,835
2018	2,004	3,965	632,488,395	295,325	593,590
2019	2,392	4,961	716,146,085	336,410	675,745
13-18	12,243		3,567,452,270	1,811,980	3,637,650

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 metrics are excluded from the table due to space limitations, but are present in the "13-18" summations.