

Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Adams	199	96.5	0.0	3.0	0.0	0.0	0.0	0.5
Allen	2,751	50.4	26.6	14.4	0.1	5.3	2.3	0.8
Bartholome	526	81.9	1.7	14.3	0.0	0.6	0.8	0.8
Benton	45	*	*	*	*	*	*	*
Blackford	102	95.1	1.0	3.9	0.0	0.0	0.0	0.0
Boone	202	92.6	3.0	2.5	0.0	1.0	0.0	1.0
Brown	64	*	*	*	*	*	*	*
Carroll	82	*	*	*	*	*	*	*
Cass	384	64.6	1.0	31.0	0.0	0.8	1.3	1.3
Clark	722	71.6	9.8	13.6	0.4	1.1	2.8	0.7
Clay	243	95.5	1.2	0.4	0.0	0.4	1.6	0.8
Clinton	269	62.5	0.4	34.9	0.4	0.0	0.7	1.1
Crawford	82	*	*	*	*	*	*	*
Daviess	196	77.6	1.0	17.9	0.0	0.0	2.0	1.5
Dearborn	330	96.7	0.9	0.3	0.3	0.0	0.6	1.2
Decatur	194	97.9	0.0	1.5	0.5	0.0	0.0	0.0
DeKalb	306	93.8	0.0	4.2	0.0	0.0	1.3	0.7
Delaware	848	83.4	11.3	2.0	0.1	0.6	2.1	0.5
Dubois	252	75.4	0.0	21.0	0.0	2.0	1.2	0.4
Elkhart	1,943	55.8	9.8	31.9	0.0	0.9	1.0	0.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Fayette	189	96.3	2.1	0.5	0.0	0.5	0.0	0.5
Floyd	476	77.3	13.0	5.0	0.2	1.9	1.7	0.8
Fountain	111	94.6	0.0	3.6	0.0	0.9	0.0	0.9
Franklin	100	98.0	0.0	1.0	0.0	0.0	0.0	1.0
Fulton	143	90.2	0.0	9.1	0.0	0.0	0.7	0.0
Gibson	200	93.5	2.5	1.5	1.0	0.5	0.5	0.5
Grant	564	77.1	13.5	5.5	0.0	0.5	2.1	1.2
Greene	164	97.0	0.0	1.8	0.0	0.0	0.6	0.6
Hamilton	682	78.3	7.2	10.4	0.1	2.5	1.3	0.1
Hancock	296	94.9	1.4	2.4	0.0	0.0	1.0	0.3
Harrison	254	90.9	0.4	5.1	0.0	0.0	2.0	1.6
Hendricks	421	85.3	6.2	3.3	0.0	3.8	0.0	1.4
Henry	300	93.3	1.0	3.0	0.3	0.3	1.3	0.7
Howard	705	85.1	8.1	3.7	0.0	0.3	1.8	1.0
Huntington	232	96.1	0.4	1.7	0.0	0.0	1.3	0.4
Jackson	352	84.1	1.1	13.9	0.0	0.0	0.0	0.9
Jasper	217	85.7	0.9	12.4	0.0	0.0	0.5	0.5
Jay	158	90.5	0.6	6.3	0.0	0.0	0.0	2.5
Jefferson	224	85.7	1.3	7.6	0.0	0.9	2.2	2.2
Jennings	284	95.4	0.7	2.1	0.7	0.4	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Johnson	730	87.8	2.1	4.9	0.0	2.6	1.2	1.4
Knox	239	95.0	0.4	0.8	0.0	0.4	1.3	2.1
Kosciusko	441	77.6	1.1	20.0	0.0	0.5	0.0	0.9
LaGrange	145	83.4	1.4	12.4	0.0	0.7	0.0	2.1
Lake	4,161	30.8	42.5	23.9	0.1	0.7	1.3	0.8
LaPorte	878	68.1	19.1	9.0	0.0	0.8	2.2	0.8
Lawrence	285	96.1	0.0	1.1	0.0	1.1	0.4	1.4
Madison	957	78.3	12.0	7.6	0.2	0.2	1.0	0.6
Marion	9,411	34.1	40.1	19.9	0.1	3.6	1.5	0.7
Marshall	373	76.4	0.3	22.0	0.0	0.3	1.1	0.0
Martin	57	*	*	*	*	*	*	*
Miami	202	90.6	2.5	3.0	0.0	0.5	3.5	0.0
Monroe	657	82.3	4.7	3.8	0.0	6.4	2.4	0.3
Montgomery	282	86.9	1.4	8.5	0.4	0.7	1.1	1.1
Morgan	423	96.9	0.7	1.7	0.0	0.0	0.2	0.5
Newton	57	*	*	*	*	*	*	*
Noble	304	79.6	0.0	18.8	0.0	1.0	0.7	0.0
Ohio	7	*	*	*	*	*	*	*
Orange	159	95.6	1.9	0.0	0.0	1.3	0.6	0.6
Owen	116	94.8	0.9	2.6	0.0	0.9	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Parke	75	*	*	*	*	*	*	*
Perry	127	99.2	0.0	0.0	0.0	0.0	0.0	0.8
Pike	74	*	*	*	*	*	*	*
Porter	786	77.6	8.1	11.6	0.0	1.0	0.1	1.5
Posey	98	*	*	*	*	*	*	*
Pulaski	70	*	*	*	*	*	*	*
Putnam	222	94.6	1.4	1.8	0.0	0.9	0.5	0.9
Randolph	190	86.8	0.0	8.4	0.5	0.5	1.6	2.1
Ripley	196	93.9	0.0	2.0	0.5	0.0	1.5	2.0
Rush	106	97.2	0.0	0.9	0.0	0.9	0.0	0.9
Scott	227	97.8	0.0	1.3	0.0	0.0	0.4	0.4
Shelby	324	87.0	1.9	9.9	0.0	0.0	0.6	0.6
Spencer	102	89.2	1.0	5.9	0.0	0.0	0.0	3.9
St Joseph	1,958	48.7	28.8	16.4	0.3	1.8	2.6	1.3
Starke	171	96.5	0.0	2.3	0.0	0.6	0.0	0.6
Steuben	275	89.5	0.7	6.2	0.0	1.1	0.4	2.2
Sullivan	158	98.7	0.0	1.3	0.0	0.0	0.0	0.0
Switzerlan	44	*	*	*	*	*	*	*
Tippecanoe	1,273	61.5	11.5	18.1	0.2	6.4	1.1	1.2
Tipton	61	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Union	48	*	*	*	*	*	*	*
Vanderburg	1,021	73.5	16.7	5.8	0.3	0.8	2.3	0.7
Vermillion	89	*	*	*	*	*	*	*
Vigo	676	83.9	10.8	2.4	0.0	0.9	1.2	0.9
Wabash	217	88.0	1.4	2.8	0.0	0.5	6.9	0.5
Warren	67	*	*	*	*	*	*	*
Warrick	246	95.5	2.4	0.4	0.0	0.4	1.2	0.0
Washington	168	98.8	0.0	0.6	0.0	0.0	0.0	0.6
Wayne	605	82.3	6.4	6.6	0.3	0.8	2.5	1.0
Wells	158	89.2	1.3	7.6	0.0	0.0	1.3	0.6
White	213	79.8	0.0	18.3	0.0	0.5	1.4	0.0
Whitley	184	97.3	0.5	1.1	0.0	0.5	0.5	0.0
Unknown Counties	307	68.7	17.3	9.1	0.0	2.0	1.0	2.0
Indiana	46,232	64.1	18.3	13.4	0.1	1.9	1.4	0.8
Nation (Prior Year)	1,162,961	41.6	22.7	29.2	1.1	3.4	1.3	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Adams	199	0.0	4.0	13.1	67.8	14.6	0.5	0.0
Allen	2,751	0.2	3.6	12.1	62.6	19.7	1.3	0.4
Bartholome	526	0.0	9.5	16.3	57.6	14.8	1.7	0.0
Benton	45	*	*	*	*	*	*	*
Blackford	102	0.0	6.9	16.7	66.7	8.8	1.0	0.0
Boone	202	0.0	5.0	15.3	67.3	11.4	1.0	0.0
Brown	64	*	*	*	*	*	*	*
Carroll	82	*	*	*	*	*	*	*
Cass	384	0.8	6.3	14.1	61.2	16.7	1.0	0.0
Clark	722	0.1	4.4	13.2	62.3	19.1	0.7	0.1
Clay	243	0.4	7.4	16.0	64.2	10.7	1.2	0.0
Clinton	269	0.0	4.5	14.1	64.7	14.9	1.9	0.0
Crawford	82	*	*	*	*	*	*	*
Daviess	196	0.0	4.6	12.8	66.8	13.3	0.0	2.6
Dearborn	330	0.6	4.8	13.0	63.0	17.3	0.9	0.3
Decatur	194	0.0	7.2	18.6	61.3	12.4	0.5	0.0
DeKalb	306	0.3	6.2	18.0	62.4	12.4	0.7	0.0
Delaware	848	0.1	5.1	13.3	65.3	15.3	0.8	0.0
Dubois	252	0.0	2.8	9.9	64.7	20.2	2.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Elkhart	1,943	0.4	4.4	13.6	59.5	20.7	1.4	0.0
Fayette	189	0.5	6.9	15.3	67.2	10.1	0.0	0.0
Floyd	476	0.4	6.3	16.2	62.4	13.9	0.6	0.2
Fountain	111	0.0	4.5	17.1	59.5	18.9	0.0	0.0
Franklin	100	0.0	9.0	12.0	62.0	16.0	1.0	0.0
Fulton	143	0.0	3.5	12.6	71.3	11.2	1.4	0.0
Gibson	200	0.0	3.0	17.5	65.0	13.5	0.5	0.5
Grant	564	1.1	5.9	15.1	64.5	12.8	0.7	0.0
Greene	164	0.6	8.5	16.5	64.6	9.8	0.0	0.0
Hamilton	682	0.3	2.2	12.3	60.4	22.3	1.5	1.0
Hancock	296	0.0	5.1	13.5	61.5	18.9	1.0	0.0
Harrison	254	0.0	3.5	14.6	65.4	15.4	0.8	0.4
Hendricks	421	0.0	4.3	14.5	60.1	19.5	1.7	0.0
Henry	300	0.3	6.0	14.0	65.3	13.3	0.7	0.3
Howard	705	0.4	5.7	15.5	62.7	14.8	0.9	0.1
Huntington	232	0.4	5.2	20.7	62.5	9.5	0.9	0.9
Jackson	352	0.0	8.0	15.1	63.6	12.5	0.9	0.0
Jasper	217	0.0	5.1	15.7	55.8	20.3	3.2	0.0
Jay	158	0.0	7.6	10.1	68.4	13.9	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Jefferson	224	0.4	7.1	14.7	62.1	14.7	0.9	0.0
Jennings	284	0.4	7.0	17.6	62.7	12.0	0.4	0.0
Johnson	730	0.1	5.1	15.1	63.0	14.8	1.9	0.0
Knox	239	0.4	2.5	16.7	66.1	14.2	0.0	0.0
Kosciusko	441	0.5	3.9	15.2	59.0	20.0	1.6	0.0
LaGrange	145	0.7	6.2	16.6	56.6	17.9	0.7	1.4
Lake	4,161	0.4	5.3	12.0	57.8	22.9	1.4	0.1
LaPorte	878	0.3	3.9	13.7	61.4	19.1	1.6	0.0
Lawrence	285	0.0	9.8	15.4	58.9	14.4	1.4	0.0
Madison	957	0.3	6.3	15.3	61.4	15.3	1.4	0.1
Marion	9,411	0.4	4.7	10.7	62.7	20.1	1.3	0.1
Marshall	373	0.3	3.8	12.1	59.8	22.5	1.3	0.3
Martin	57	*	*	*	*	*	*	*
Miami	202	0.5	5.0	16.3	61.4	16.3	0.0	0.5
Monroe	657	0.0	2.1	13.1	63.6	19.3	1.2	0.6
Montgomery	282	0.0	7.8	14.2	60.6	16.0	1.4	0.0
Morgan	423	0.0	5.0	11.3	67.4	15.1	0.9	0.2
Newton	57	*	*	*	*	*	*	*
Noble	304	0.0	6.3	14.1	63.5	15.1	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Ohio	7	*	*	*	*	*	*	*
Orange	159	0.6	3.8	14.5	60.4	20.1	0.6	0.0
Owen	116	0.0	2.6	19.8	64.7	12.9	0.0	0.0
Parke	75	*	*	*	*	*	*	*
Perry	127	0.8	10.2	17.3	63.0	7.9	0.8	0.0
Pike	74	*	*	*	*	*	*	*
Porter	786	0.1	4.8	11.6	62.3	19.2	1.7	0.3
Posey	98	*	*	*	*	*	*	*
Pulaski	70	*	*	*	*	*	*	*
Putnam	222	0.5	5.9	13.5	62.6	16.7	0.9	0.0
Randolph	190	0.5	5.8	15.3	61.6	15.3	1.1	0.5
Ripley	196	0.0	4.1	13.8	68.4	12.8	1.0	0.0
Rush	106	0.0	10.4	19.8	61.3	7.5	0.9	0.0
Scott	227	0.4	7.0	15.4	67.8	7.9	1.3	0.0
Shelby	324	0.3	8.3	12.7	64.8	12.7	1.2	0.0
Spencer	102	0.0	6.9	13.7	54.9	21.6	2.9	0.0
St Joseph	1,958	0.1	6.7	13.1	58.1	20.9	1.1	0.0
Starke	171	1.2	2.9	10.5	64.9	18.1	1.8	0.6
Steuben	275	0.0	5.5	19.3	63.6	11.3	0.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sullivan	158	0.0	5.1	22.8	59.5	12.7	0.0	0.0
Switzerlan	44	*	*	*	*	*	*	*
Tippecanoe	1,273	0.1	4.5	12.8	60.5	20.5	1.6	0.1
Tipton	61	*	*	*	*	*	*	*
Union	48	*	*	*	*	*	*	*
Vanderburg	1,021	0.2	6.8	12.5	64.9	14.7	0.8	0.1
Vermillion	89	*	*	*	*	*	*	*
Vigo	676	0.3	8.4	16.1	63.2	11.2	0.7	0.0
Wabash	217	0.0	5.5	14.3	66.4	11.5	1.4	0.9
Warren	67	*	*	*	*	*	*	*
Warrick	246	0.0	7.3	15.4	65.9	10.6	0.0	0.8
Washington	168	0.0	5.4	19.0	62.5	11.9	0.6	0.6
Wayne	605	0.0	3.5	14.7	64.6	16.7	0.5	0.0
Wells	158	0.6	1.9	20.3	64.6	12.7	0.0	0.0
White	213	0.5	6.1	14.6	59.2	18.3	1.4	0.0
Whitley	184	0.0	3.3	11.4	68.5	15.8	1.1	0.0
Unknown Counties	307	0.3	6.5	11.7	65.8	14.3	1.3	0.0
Indiana	46,232	0.3	5.2	13.2	62.1	17.9	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Nation (Prior Year)	1,162,961	0.3	5.3	11.5	59.4	21.7	1.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams	167	6.0 (46)	54.5 (37)	161	19.9 (45)	52.8 (29)	40	* *	35	* *	195	1.0 (9)	195	3.6(42)
Allen	2,241	4.4 (16)	54.4 (35)	2,143	21.2 (58)	49.7 (11)	377	41.9 (14)	386	32.4 (17)	2,675	2.5 (25)	2,675	3.9(45)
Bartholome	424	7.3 (66)	53.8 (31)	412	18.9 (38)	54.4 (44)	69	* *	70	* *	516	4.7 (62)	516	7.4(67)
Benton	36	* *	* *	34	* *	* *	7	* *	8	* *	43	* *	43	* *
Blackford	80	* *	* *	76	* *	* *	22	* *	20	* *	99	* *	99	* *
Boone	158	5.1 (28)	50.0 (9)	155	14.8 (4)	54.2 (43)	30	* *	36	* *	198	2.0 (21)	198	0.5(3)
Brown	50	* *	* *	48	* *	* *	8	* *	18	* *	62	* *	62	* *
Carroll	66	* *	* *	63	* *	* *	10	* *	16	* *	81	* *	81	* *
Cass	326	4.3 (15)	55.5 (43)	311	22.5 (65)	47.9 (4)	61	* *	36	* *	372	3.0 (37)	372	0.8(7)
Clark	577	5.7 (39)	58.8 (62)	557	18.9 (38)	55.3 (49)	110	35.5 (7)	112	33.9(20)	704	4.4 (58)	704	1.6(18)
Clay	207	6.3 (54)	57.5 (55)	199	22.1 (61)	48.2 (6)	28	* *	53	* *	237	3.4 (45)	237	2.5(31)
Clinton	220	3.2 (4)	63.2 (68)	218	17.4 (23)	49.5 (10)	51	* *	38	* *	267	3.4 (45)	267	3.0(37)
Crawford	64	* *	* *	62	* *	* *	13	* *	13	* *	80	* *	80	* *
Daviess	164	2.4 (1)	62.8 (67)	153	17.0 (20)	52.9 (30)	32	* *	111	40.5(23)	187	2.1 (23)	187	5.3(55)
Dearborn	249	6.0 (46)	51.0 (16)	238	16.8 (14)	61.8 (69)	46	* *	72	* *	321	1.6 (15)	321	1.6(18)
Decatur	151	6.0 (46)	53.6 (29)	147	15.0 (5)	50.3 (15)	25	* *	32	* *	190	3.2 (41)	190	6.8(61)
DeKalb	253	6.7 (56)	51.4 (17)	235	21.3 (59)	56.2 (53)	40	* *	42	* *	295	1.7 (16)	295	1.4(17)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Delaware	670	5.5 (35)	56.4 (50)	640	19.1 (41)	56.6 (57)	116	40.5 (12)	113	38.1(22)	827	2.8 (32)	827	3.3(41)
Dubois	191	5.8 (41)	60.7 (65)	185	16.8 (14)	51.9 (24)	41	* *	36	* *	249	4.4 (58)	249	2.8(34)
Elkhart	1,656	4.0 (11)	57.5 (55)	1,599	19.9 (45)	51.2 (21)	191	27.2 (1)	247	17.4 (2)	1,893	2.7 (30)	1,893	2.1(28)
Fayette	146	6.8 (58)	55.5 (43)	140	18.6 (36)	50.7 (19)	22	* *	33	* *	183	1.1 (10)	183	1.1(11)
Floyd	377	5.0 (26)	53.6 (29)	361	20.2 (49)	52.9 (30)	74	* *	77	* *	461	6.3 (71)	461	9.8(71)
Fountain	92	* *	* *	87	* *	* *	14	* *	23	* *	106	1.9 (19)	106	7.5(68)
Franklin	81	* *	* *	80	* *	* *	12	* *	14	* *	99	* *	99	* *
Fulton	117	5.1 (28)	60.7 (65)	104	16.3 (12)	61.5 (68)	16	* *	36	* *	131	0.0 (1)	131	0.8(7)
Gibson	150	5.3 (32)	50.0 (9)	137	22.6 (66)	53.3 (35)	35	* *	31	* *	187	1.1 (10)	187	1.1(11)
Grant	450	5.3 (32)	49.8 (8)	422	19.0 (40)	51.7 (23)	87	* *	238	24.8(10)	541	3.3 (43)	541	7.2(63)
Greene	129	6.2 (52)	54.3 (34)	123	20.3 (51)	56.1 (52)	26	* *	44	* *	161	1.9 (19)	161	0.6(5)
Hamilton	527	4.9 (22)	46.3 (1)	505	18.4 (31)	53.9 (41)	130	37.7 (10)	158	27.2(14)	663	5.1 (65)	663	3.2(39)
Hancock	226	4.9 (22)	53.1 (24)	216	15.7 (7)	53.2 (32)	54	* *	68	* *	288	3.1 (40)	288	1.0(10)
Harrison	200	5.5 (35)	50.0 (9)	190	21.1 (56)	50.0 (12)	26	* *	37	* *	245	6.5 (72)	245	5.7(58)
Hendricks	310	3.2 (4)	52.9 (23)	261	11.5 (1)	58.2 (64)	88	* *	217	47.9(24)	373	2.7 (30)	373	1.6(18)
Henry	253	7.1 (62)	48.2 (2)	249	16.9 (17)	56.6 (57)	39	* *	26	* *	297	3.0 (37)	297	0.3(2)
Howard	570	5.6 (37)	55.3 (42)	553	17.2 (21)	50.6 (17)	82	* *	123	23.6 (9)	690	1.2 (12)	690	1.3(15)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Huntington	196	7.1 (62)	49.0 (5)	187	20.9 (55)	48.7 (9)	25	* *	27	* *	224	4.5 (60)	224	5.8(59)
Jackson	288	5.9 (44)	55.9 (47)	276	18.8 (37)	53.3 (35)	49	* *	75	* *	344	3.8 (49)	344	5.2(53)
Jasper	171	4.1 (13)	54.4 (35)	162	17.3 (22)	56.2 (53)	14	* *	38	* *	209	0.0 (1)	209	0.0(1)
Jay	113	6.2 (52)	58.4 (59)	111	25.2 (67)	50.5 (16)	22	* *	38	* *	157	2.5 (25)	157	2.5(31)
Jefferson	171	7.0 (61)	55.0 (41)	162	18.5 (35)	51.9 (24)	27	* *	36	* *	219	4.1 (54)	219	5.5(56)
Jennings	243	5.8 (41)	54.7 (39)	229	18.3 (30)	53.7 (37)	21	* *	48	* *	274	2.6 (29)	274	2.2(29)
Johnson	526	6.8 (58)	50.8 (14)	506	15.6 (6)	56.9 (59)	133	42.1 (15)	144	19.4 (5)	712	3.8 (49)	712	8.0(69)
Knox	194	8.2 (67)	51.5 (19)	168	16.1 (11)	58.3 (65)	31	* *	125	20.0 (6)	215	5.6 (68)	215	4.2(47)
Kosciusko	342	5.6 (37)	53.5 (28)	325	20.6 (54)	48.3 (7)	75	* *	106	16.0 (1)	425	4.2 (56)	425	2.8(34)
LaGrange	111	7.2 (65)	51.4 (17)	106	25.5 (68)	50.0 (12)	26	* *	35	* *	139	4.3 (57)	139	4.3(49)
Lake	2,987	4.2 (14)	59.3 (63)	2,705	18.2 (28)	55.6 (51)	682	41.5 (13)	564	29.4(16)	3,901	2.9 (35)	3,901	4.2(47)
LaPorte	696	5.9 (44)	56.2 (49)	650	16.9 (17)	53.7 (37)	147	35.4 (6)	167	25.7(11)	839	2.4 (24)	839	1.7(22)
Lawrence	237	6.8 (58)	54.9 (40)	233	16.3 (12)	53.2 (32)	33	* *	51	* *	283	5.3 (67)	283	8.1(70)
Madison	735	4.5 (19)	56.5 (52)	702	18.4 (31)	52.4 (27)	152	48.7 (16)	165	36.4(21)	931	1.5 (14)	931	1.7(22)
Marion	6,641	4.4 (16)	58.2 (58)	6,156	20.2 (49)	51.6 (22)	1,384	36.9 (8)	1,391	26.2(12)	8,973	4.5 (60)	8,973	5.1(52)
Marshall	296	2.4 (1)	60.1 (64)	279	15.8 (9)	54.5 (45)	60	* *	63	* *	362	4.7 (62)	362	5.5(56)
Martin	47	* *	* *	46	* *	* *	4	* *	1	* *	56	* *	56	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	166	6.0 (46)	50.0 (9)	160	17.5 (25)	55.0 (48)	28	* *	115	28.7 (15)	197	2.5 (25)	197	6.1 (60)
Monroe	526	4.9 (22)	53.8 (31)	484	18.4 (31)	53.7 (37)	103	27.2 (1)	145	32.4 (17)	615	3.3 (43)	615	0.8 (7)
Montgomery	225	4.4 (16)	53.3 (25)	214	16.8 (14)	57.9 (62)	48	* *	53	* *	270	4.1 (54)	270	1.1 (11)
Morgan	299	6.0 (46)	58.5 (60)	279	13.6 (3)	58.1 (63)	60	* *	65	* *	407	2.0 (21)	407	1.7 (22)
Newton	43	* *	* *	43	* *	* *	7	* *	10	* *	57	* *	57	* *
Noble	250	3.2 (4)	57.6 (57)	235	19.6 (42)	54.9 (47)	55	* *	49	* *	291	0.0 (1)	291	1.7 (22)
Ohio	4	* *	* *	4	* *	* *	1	* *	0	* *	7	* *	7	* *
Orange	122	3.3 (8)	65.6 (70)	116	22.4 (64)	47.4 (3)	16	* *	27	* *	155	5.2 (66)	155	11.6 (74)
Owen	99	* *	* *	96	* *	* *	9	* *	25	* *	113	10.6 (73)	113	11.5 (73)
Parke	59	* *	* *	58	* *	* *	8	* *	19	* *	75	* *	75	* *
Perry	97	* *	* *	93	* *	* *	16	* *	22	* *	124	3.2 (41)	124	7.3 (65)
Pike	61	* *	* *	56	* *	* *	10	* *	6	* *	69	* *	69	* *
Porter	598	4.0 (11)	51.8 (20)	562	18.1 (27)	53.9 (41)	113	29.2 (3)	152	27.0 (13)	751	2.5 (25)	751	2.9 (36)
Posey	75	* *	* *	70	* *	* *	15	* *	18	* *	93	* *	93	* *
Pulaski	54	* *	* *	52	* *	* *	7	* *	7	* *	68	* *	68	* *
Putnam	188	3.2 (4)	48.4 (3)	184	17.4 (23)	57.1 (61)	20	* *	47	* *	219	11.0 (74)	219	7.3 (65)
Randolph	158	8.2 (67)	55.7 (46)	148	27.7 (69)	47.3 (2)	34	* *	30	* *	186	0.5 (5)	186	1.6 (18)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Ripley	145	3.4 (9)	55.9 (47)	137	12.4 (2)	52.6 (28)	22	* *	41	* *	188	5.9 (70)	188	6.9(62)
Rush	89	* *	* *	88	* *	* *	13	* *	16	* *	105	4.8 (64)	105	3.8(44)
Scott	191	9.9 (70)	53.9 (33)	189	22.2 (63)	54.5 (45)	14	* *	25	* *	225	3.6 (47)	225	4.0(46)
Shelby	256	6.3 (54)	50.0 (9)	247	18.2 (28)	48.6 (8)	48	* *	90	* *	317	3.8 (49)	317	1.9(27)
Spencer	82	* *	* *	80	* *	* *	21	* *	12	* *	102	2.9 (35)	102	4.9(51)
St Joseph	1,561	4.9 (22)	55.5 (43)	1,489	20.3 (51)	50.2 (14)	290	39.0 (11)	417	32.9(19)	1,895	1.7 (16)	1,895	2.2(29)
Starke	150	6.7 (56)	49.3 (7)	146	19.9 (45)	56.2 (53)	20	* *	15	* *	167	3.0 (37)	167	1.2(14)
Steuben	234	5.1 (28)	56.4 (50)	227	19.8 (44)	53.7 (37)	26	* *	31	* *	269	0.7 (7)	269	0.7(6)
Sullivan	139	5.8 (41)	63.3 (69)	133	20.3 (51)	51.1 (20)	20	* *	53	* *	153	0.7 (7)	153	2.6(33)
Switzerlan	30	* *	* *	29	* *	* *	5	* *	13	* *	42	* *	42	* *
Tippecanoe	1,045	4.6 (20)	52.2 (21)	984	20.1 (48)	48.0 (5)	172	37.2 (9)	156	18.6 (3)	1,227	2.8 (32)	1,227	1.3(15)
Tipton	46	* *	* *	42	* *	* *	9	* *	8	* *	57	* *	57	* *
Union	35	* *	* *	34	* *	* *	4	* *	9	* *	47	* *	47	* *
Vanderburg	868	5.3 (32)	57.3 (54)	757	18.4 (31)	58.9 (66)	175	33.7 (5)	421	21.1 (7)	917	2.8 (32)	917	3.2(39)
Vermillion	73	* *	* *	71	* *	* *	13	* *	7	* *	87	* *	87	* *
Vigo	557	5.2 (31)	50.8 (14)	475	15.8 (9)	59.2 (67)	102	32.4 (4)	340	22.4 (8)	600	1.8 (18)	600	3.7(43)
Wabash	182	7.1 (62)	53.3 (25)	172	15.7 (7)	52.3 (26)	33	* *	128	18.8 (4)	208	3.8 (49)	208	7.2(63)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
Warren	50	* *	* *	50	* *	* *	7	* *	21	* *	67	* *	67	* *
Warrick	205	3.4 (9)	52.7 (22)	193	17.6 (26)	56.5 (56)	36	* *	32	* *	236	0.4 (4)	236	3.0(37)
Washington	145	4.8 (21)	58.6 (61)	138	19.6 (42)	47.1 (1)	14	* *	20	* *	163	3.7 (48)	163	9.8(71)
Wayne	489	5.7 (39)	56.9 (53)	462	22.1 (61)	50.6 (17)	68	* *	88	* *	586	5.8 (69)	586	4.3(49)
Wells	119	5.0 (26)	54.6 (38)	114	21.1 (56)	55.3 (49)	30	* *	26	* *	155	1.3 (13)	155	5.2(53)
White	177	2.8 (3)	49.2 (6)	173	21.4 (60)	53.2 (32)	17	* *	26	* *	208	0.5 (5)	208	0.5(3)
Whitley	150	6.0 (46)	53.3 (25)	142	16.9 (17)	57.0 (60)	28	* *	47	* *	178	3.9 (53)	178	1.7(22)
Unknown Counties	295	8.8 (69)	48.5 (4)	1	* *	* *	78	* *	2	* *	3	* *	3	* *
Indiana	35,909	5.0	55.4	33,494	19.1	52.8	6,722	36.2	8,527	26.8	44,078	3.3	44,078	3.7
Nation (Prior Year)	1,106,370	4.5	53.4	835,927	21.5	48.0	164,201	33.9	346,981	28.1	419,956	5.2	467,675	6.3

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams	132	53.0 (48)	42.4	(48)	170	48.2 (24)	167	25.1 (2)	167	93.4 (7)
Allen	1,715	58.7 (39)	36.8	(40)	2,241	45.9 (32)	2,242	17.2 (33)	2,243	89.2 (30)
Bartholome	354	44.6 (55)	45.2	(53)	425	44.5 (39)	425	18.6 (23)	426	90.1 (24)
Benton	26	* *	* *		35	* *	36	* *	36	* *
Blackford	63	* *	* *		81	* *	81	* *	81	* *
Boone	122	58.2 (41)	37.7	(42)	159	49.1 (21)	159	6.3 (70)	159	90.6 (18)
Brown	42	* *	* *		50	* *	51	* *	51	* *
Carroll	50	* *	* *		68	* *	66	* *	66	* *
Cass	256	58.6 (40)	34.0	(35)	327	45.0 (34)	326	11.0 (63)	326	82.8 (59)
Clark	431	65.7 (28)	29.5	(29)	581	41.3 (48)	578	19.6 (19)	578	92.0 (13)
Clay	141	53.2 (47)	41.1	(47)	207	54.6 (9)	206	18.9 (21)	207	89.9 (26)
Clinton	168	63.1 (33)	29.8	(30)	220	38.6 (60)	221	20.8 (14)	221	92.8 (8)
Crawford	50	* *	* *		65	* *	64	* *	64	* *
Daviess	110	77.3 (4)	17.3	(4)	164	46.3 (30)	164	12.2 (56)	164	81.1 (66)
Dearborn	144	75.0 (6)	20.1	(8)	252	36.5 (62)	249	18.1 (27)	249	90.4 (19)
Decatur	119	66.4 (26)	30.3	(31)	156	42.3 (46)	150	14.0 (51)	151	84.1 (56)
DeKalb	190	67.4 (23)	28.4	(27)	250	50.4 (19)	255	17.6 (31)	254	89.8 (27)
Delaware	462	74.5 (8)	21.4	(13)	673	47.5 (26)	667	18.0 (29)	671	90.3 (20)
Dubois	124	75.0 (6)	21.0	(11)	193	34.2 (67)	191	22.5 (7)	191	86.9 (46)
Elkhart	1,424	40.2 (57)	56.1	(59)	1,654	57.3 (6)	1,658	21.2 (11)	1,658	81.4 (65)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Fayette	90	* *	* *	* *	145	51.7 (15)	146	11.6 (60)	146	86.3 (49)
Floyd	267	65.9 (27)	25.8 (23)		385	40.3 (56)	378	20.4 (17)	378	89.2 (30)
Fountain	80	* *	* *	* *	93	* *	92	* *	92	* *
Franklin	58	* *	* *	* *	82	* *	81	* *	81	* *
Fulton	74	* *	* *	* *	120	46.7 (29)	117	15.4 (47)	117	88.0 (41)
Gibson	94	* *	* *	* *	154	39.6 (58)	151	20.5 (16)	151	94.7 (2)
Grant	304	68.8 (20)	23.7 (16)		451	41.2 (50)	451	16.4 (40)	451	88.5 (37)
Greene	71	* *	* *	* *	131	45.0 (34)	100	11.0 (63)	127	93.7 (5)
Hamilton	362	74.0 (10)	21.3 (12)		533	37.5 (61)	526	21.5 (9)	528	92.2 (11)
Hancock	181	64.1 (31)	31.5 (33)		226	42.5 (45)	228	20.6 (15)	228	90.8 (17)
Harrison	137	50.4 (52)	38.7 (44)		202	47.5 (26)	201	24.9 (4)	201	96.0 (1)
Hendricks	194	82.0 (2)	11.3 (1)		314	29.0 (69)	306	22.5 (7)	311	92.6 (9)
Henry	215	61.9 (36)	36.3 (39)		254	52.4 (13)	254	15.0 (48)	254	81.5 (64)
Howard	423	62.4 (35)	33.1 (34)		574	46.0 (31)	571	9.6 (67)	572	87.9 (42)
Huntington	149	44.3 (56)	54.4 (57)		195	61.5 (3)	196	21.4 (10)	197	85.3 (55)
Jackson	212	74.5 (8)	20.8 (9)		291	44.7 (37)	288	13.2 (55)	289	92.0 (13)
Jasper	121	62.8 (34)	35.5 (37)		170	48.2 (24)	171	14.0 (51)	171	85.4 (54)
Jay	83	* *	* *	* *	112	49.1 (21)	113	11.5 (61)	113	89.4 (29)
Jefferson	141	51.8 (50)	44.0 (51)		176	54.0 (12)	172	16.9 (37)	172	83.7 (57)
Jennings	195	29.2 (60)	67.7 (60)		239	67.4 (1)	243	16.0 (41)	243	74.9 (69)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	418	63.4 (32)	29.4 (28)		532	33.3 (68)	508	16.5 (38)	526	90.3 (20)
Knox	124	78.2 (3)	19.4 (6)		200	41.0 (52)	197	9.1 (68)	197	88.8 (33)
Kosciusko	248	76.2 (5)	16.9 (3)		340	45.0 (34)	343	15.7 (44)	342	86.5 (47)
LaGrange	76	* *	* *		114	43.9 (42)	111	18.9 (21)	111	86.5 (47)
Lake	1,961	70.8 (15)	19.8 (7)		3,012	34.5 (66)	2,852	15.7 (44)	2,975	88.2 (39)
LaPorte	552	56.2 (45)	36.8 (40)		697	40.9 (54)	665	20.2 (18)	688	93.9 (4)
Lawrence	177	59.3 (38)	36.2 (38)		234	54.7 (8)	237	10.1 (66)	238	81.9 (62)
Madison	563	64.8 (29)	23.4 (15)		739	41.1 (51)	737	14.9 (50)	737	90.1 (24)
Marion	4,976	67.1 (24)	24.4 (18)		6,648	35.7 (64)	6,605	11.7 (59)	6,659	87.5 (44)
Marshall	241	60.6 (37)	35.3 (36)		296	40.2 (57)	296	19.3 (20)	297	88.2 (39)
Martin	32	* *	* *		45	* *	47	* *	47	* *
Miami	111	67.6 (22)	30.6 (32)		168	51.2 (17)	166	17.5 (32)	166	89.2 (30)
Monroe	349	69.6 (16)	25.8 (23)		522	44.4 (40)	380	23.4 (5)	508	92.1 (12)
Montgomery	185	52.4 (49)	43.2 (50)		222	44.6 (38)	225	16.0 (41)	225	90.2 (23)
Morgan	224	69.2 (19)	24.1 (17)		303	35.0 (65)	300	21.0 (12)	300	90.3 (20)
Newton	33	* *	* *		40	* *	43	* *	43	* *
Noble	178	64.6 (30)	25.3 (22)		248	44.4 (40)	250	18.4 (25)	250	88.8 (33)
Ohio	4	* *	* *		4	* *	4	* *	4	* *
Orange	88	* *	* *		120	54.2 (11)	121	16.5 (38)	122	82.8 (59)
Owen	73	* *	* *		98	* *	99	* *	99	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prenpregnancy (4)		During Preganancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Parke	40	* *	* *	* *	59	* *	59	* *	58	* *
Perry	83	* *	* *	* *	99	* *	98	* *	98	* *
Pike	36	* *	* *	* *	62	* *	61	* *	61	* *
Porter	416	67.8 (21)	27.6 (25)		599	43.1 (43)	598	20.9 (13)	598	89.5 (28)
Posey	56	* *	* *	* *	76	* *	75	* *	75	* *
Pulaski	32	* *	* *	* *	54	* *	54	* *	54	* *
Putnam	144	47.9 (53)	48.6 (54)		187	52.4 (13)	188	25.0 (3)	187	83.4 (58)
Randolph	106	82.1 (1)	13.2 (2)		157	42.7 (44)	157	10.8 (65)	158	93.7 (5)
Ripley	106	67.0 (25)	28.3 (26)		146	39.0 (59)	147	18.4 (25)	147	82.3 (61)
Rush	63	* *	* *	* *	86	* *	88	* *	89	* *
Scott	169	56.8 (43)	40.8 (46)		192	58.9 (4)	192	15.6 (46)	192	81.8 (63)
Shelby	170	69.4 (17)	24.7 (19)		252	41.3 (48)	257	8.9 (69)	256	85.9 (53)
Spencer	66	* *	* *	* *	81	* *	82	* *	82	* *
St Joseph	1,214	69.4 (17)	25.1 (20)		1,577	40.5 (55)	1,563	17.1 (34)	1,563	92.5 (10)
Starke	126	54.0 (46)	44.4 (52)		149	55.0 (7)	150	16.0 (41)	150	79.3 (68)
Steuben	177	46.3 (54)	52.0 (56)		234	62.4 (2)	234	15.0 (48)	234	79.9 (67)
Sullivan	74	* *	* *	* *	138	50.7 (18)	138	18.1 (27)	139	87.1 (45)
Switzerlan	25	* *	* *	* *	31	* *	31	* *	31	* *
Tippecanoe	833	50.8 (51)	42.7 (49)		1,037	51.7 (15)	982	23.1 (6)	1,034	86.0 (51)
Tipton	26	* *	* *	* *	45	* *	46	* *	46	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before preanancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Union	22	* *	* *	* *	34	* *	35	* *	35	* *
Vanderburg	577	71.9 (13)	20.8	(9)	883	36.4 (63)	869	13.7 (53)	868	90.9 (16)
Vermillion	47	* *	* *	* *	73	* *	73	* *	73	* *
Vigo	312	73.4 (12)	18.6	(5)	559	41.5 (47)	559	11.3 (62)	560	88.6 (35)
Wabash	127	70.9 (14)	25.2	(21)	183	47.0 (28)	182	17.0 (35)	182	91.8 (15)
Warren	44	* *	* *	* *	50	* *	50	* *	50	* *
Warrick	141	56.7 (44)	38.3	(43)	206	45.6 (33)	205	12.2 (56)	206	87.9 (42)
Washington	118	35.6 (59)	55.1	(58)	140	57.9 (5)	145	17.9 (30)	145	86.2 (50)
Wayne	415	38.8 (58)	51.6	(55)	487	54.6 (9)	425	13.4 (54)	471	74.3 (70)
Wells	90	* *	* *	* *	122	41.0 (52)	119	18.5 (24)	120	94.2 (3)
White	126	73.8 (11)	22.2	(14)	176	48.3 (23)	178	11.8 (58)	178	86.0 (51)
Whitley	112	58.0 (42)	40.2	(45)	149	50.3 (20)	150	28.0 (1)	149	88.6 (35)
Unknown Counties	1	* *	* *	* *	305	16.7 (70)	294	17.0 (35)	296	88.5 (37)
Indiana	26,281	62.8	30.8		36,028	42.6	35,411	16.1	35,904	88.0
Nation (Prior Year)	429,770	83.1	4.1		1,074,863	36.0	380,154	19.2	418,448	85.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams	199	31.2 (17)	195	19.5 (14)	61	* *	167	46.1 (34)	193	41.5 (36)
Allen	2,751	27.3 (11)	2,675	17.5 (7)	717	31.9 (20)	2,235	36.3 (19)	2,633	32.9 (22)
Bartholome	526	37.5 (40)	516	26.6 (41)	192	30.2 (26)	426	54.7 (57)	510	50.4 (58)
Benton	45	* *	43	* *	16	* *	36	* *	43	* *
Blackford	102	38.2 (49)	99	* *	38	* *	81	* *	99	* *
Boone	202	41.1 (59)	198	33.3 (65)	81	* *	159	51.6 (49)	195	51.8 (63)
Brown	64	* *	62	* *	25	* *	51	* *	61	* *
Carroll	82	* *	81	* *	23	* *	66	* *	80	* *
Cass	384	26.3 (8)	372	18.0 (8)	95	* *	326	31.6 (12)	366	24.9 (17)
Clark	722	31.6 (19)	704	22.4 (22)	216	33.8 (15)	578	38.9 (23)	699	35.6 (25)
Clay	243	37.9 (44)	237	26.2 (39)	89	* *	207	59.4 (67)	234	57.7 (70)
Clinton	269	26.0 (6)	267	18.4 (10)	70	* *	221	20.8 (3)	261	6.1 (2)
Crawford	82	* *	80	* *	29	* *	64	* *	78	* *
Daviess	196	37.8 (42)	187	23.5 (28)	69	* *	164	48.8 (42)	184	46.2 (47)
Dearborn	330	42.7 (64)	321	29.0 (58)	136	41.2 (6)	249	48.2 (40)	312	43.6 (42)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Decatur	194	38.1 (48)	190	28.9 (57)	71	* *	151	58.9 (66)	188	55.9 (69)
DeKalb	306	33.0 (23)	295	27.5 (50)	99	* *	255	58.4 (65)	292	49.3 (57)
Delaware	848	33.8 (26)	827	25.6 (32)	280	23.6 (34)	670	30.4 (11)	812	14.2 (7)
Dubois	252	27.0 (10)	249	21.3 (20)	66	* *	186	43.0 (26)	246	40.7 (35)
Elkhart	1,943	24.2 (5)	1,893	14.7 (3)	452	32.1 (19)	1,654	19.3 (2)	1,875	8.0 (4)
Fayette	189	36.5 (38)	183	25.7 (33)	66	* *	146	61.6 (70)	182	54.4 (66)
Floyd	475	38.3 (50)	461	26.0 (36)	174	37.9 (8)	377	43.5 (28)	456	41.9 (37)
Fountain	111	47.7 (75)	106	34.0 (67)	52	* *	91	* *	105	60.0 (71)
Franklin	100	38.0 (46)	99	* *	38	* *	81	* *	97	* *
Fulton	143	36.4 (37)	131	26.0 (36)	49	* *	117	47.9 (37)	131	42.0 (38)
Gibson	200	43.5 (66)	187	39.0 (72)	79	* *	151	57.0 (62)	186	55.4 (68)
Grant	564	33.0 (23)	541	27.0 (46)	176	26.7 (30)	451	35.5 (18)	535	9.7 (5)
Greene	163	39.3 (54)	161	26.1 (38)	61	* *	129	61.2 (69)	159	50.9 (59)
Hamilton	682	22.0 (3)	663	12.2 (2)	143	47.6 (3)	527	27.7 (7)	653	19.9 (12)
Hancock	296	29.4 (16)	288	19.1 (12)	85	* *	226	24.8 (6)	286	14.0 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harrison	254	43.3 (65)	245	29.8 (61)	106	34.9 (11)	201	49.8 (45)	240	34.2 (24)
Hendricks	421	27.6 (12)	373	15.3 (6)	94	* *	311	39.9 (24)	368	38.9 (32)
Henry	300	47.7 (75)	297	33.7 (66)	140	26.4 (31)	253	53.4 (54)	294	48.0 (51)
Howard	705	35.6 (33)	690	26.7 (42)	245	27.3 (29)	572	47.9 (37)	680	36.8 (26)
Huntington	232	46.1 (72)	224	28.6 (55)	101	31.7 (24)	197	52.8 (52)	221	51.6 (62)
Jackson	352	36.1 (34)	344	22.4 (22)	123	39.0 (7)	289	38.8 (22)	341	33.1 (23)
Jasper	217	35.5 (32)	209	28.7 (56)	74	* *	171	60.8 (68)	208	46.6 (50)
Jay	158	32.9 (22)	157	26.8 (44)	51	* *	113	48.7 (41)	153	37.3 (28)
Jefferson	224	44.6 (68)	219	36.5 (70)	96	* *	172	58.1 (63)	213	43.7 (43)
Jennings	284	39.8 (58)	274	31.4 (64)	108	18.5 (35)	243	56.8 (61)	271	48.7 (53)
Johnson	730	27.8 (14)	712	21.2 (19)	195	32.8 (17)	527	42.7 (25)	699	29.9 (20)
Knox	239	39.7 (57)	215	28.4 (53)	87	* *	196	46.4 (35)	210	45.2 (45)
Kosciusko	441	26.1 (7)	425	20.9 (17)	110	25.5 (32)	343	38.5 (20)	420	23.3 (15)
LaGrange	144	26.4 (9)	139	20.9 (17)	35	* *	111	43.2 (27)	134	39.6 (34)
Lake	4,161	18.4 (1)	3,901	11.0 (1)	697	49.9 (1)	2,988	28.5 (8)	3,857	24.1 (16)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
LaPorte	878	32.8 (21)	839	24.7 (30)	267	33.0 (16)	694	43.5 (28)	824	38.8 (31)
Lawrence	285	41.4 (61)	283	29.0 (58)	117	35.0 (10)	238	45.0 (32)	280	42.9 (39)
Madison	957	36.3 (35)	931	27.1 (48)	331	30.5 (25)	737	51.3 (48)	922	46.3 (49)
Marion	9,411	18.7 (2)	8,973	14.8 (4)	1,621	37.1 (9)	6,242	29.5 (9)	8,685	28.6 (19)
Marshall	373	28.7 (15)	362	19.9 (15)	100	34.0 (14)	296	17.6 (1)	361	5.8 (1)
Martin	57	* *	56	* *	25	* *	47	* *	55	* *
Miami	202	46.0 (71)	197	25.4 (31)	89	* *	166	56.6 (59)	197	46.2 (47)
Monroe	657	33.6 (25)	615	22.9 (26)	201	31.8 (21)	526	33.7 (16)	612	22.1 (13)
Montgomery	282	37.9 (44)	270	27.0 (46)	101	34.7 (12)	224	56.7 (60)	268	43.3 (41)
Morgan	423	34.5 (28)	407	30.2 (62)	136	29.4 (27)	298	51.0 (46)	403	48.4 (52)
Newton	57	* *	57	* *	12	* *	43	* *	57	* *
Noble	304	34.5 (28)	291	27.5 (50)	99	* *	250	47.6 (36)	290	36.9 (27)
Ohio	7	* *	7	* *	0	* *	4	* *	7	* *
Orange	159	35.2 (31)	155	28.4 (53)	52	* *	122	32.8 (14)	152	19.1 (10)
Owen	116	50.0 (77)	113	39.8 (74)	57	* *	99	* *	113	51.3 (61)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Parke	75	* *	75	* *	33	* *	58	* *	75	* *
Perry	127	46.5 (73)	124	39.5 (73)	56	* *	98	* *	123	61.8 (73)
Pike	74	* *	69	* *	28	* *	61	* *	67	* *
Porter	785	33.9 (27)	751	18.9 (11)	254	48.0 (2)	596	32.9 (15)	734	22.6 (14)
Posey	98	* *	93	* *	41	* *	75	* *	93	* *
Pulaski	70	* *	68	* *	16	* *	54	* *	67	* *
Putnam	222	44.6 (68)	219	28.3 (52)	98	* *	188	51.1 (47)	217	48.8 (54)
Randolph	190	32.1 (20)	186	24.2 (29)	60	* *	158	48.1 (39)	177	37.9 (30)
Ripley	196	39.3 (54)	188	35.1 (68)	70	* *	146	58.2 (64)	182	53.8 (65)
Rush	106	47.2 (74)	105	30.5 (63)	49	* *	89	* *	103	55.3 (67)
Scott	227	41.9 (63)	225	35.6 (69)	93	* *	192	55.2 (58)	224	52.7 (64)
Shelby	324	37.0 (39)	317	25.9 (35)	113	32.7 (18)	257	52.5 (51)	313	51.1 (60)
Spencer	102	36.3 (35)	102	22.5 (24)	37	* *	81	* *	97	* *
St Joseph	1,958	27.7 (13)	1,895	14.9 (5)	516	43.8 (5)	1,560	24.7 (5)	1,868	16.0 (9)
Starke	171	35.1 (30)	167	26.9 (45)	59	* *	150	46.0 (33)	166	48.8 (54)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Steuben	275	39.6 (56)	269	22.7 (25)	107	31.8 (21)	233	54.1 (55)	263	45.6 (46)
Sullivan	158	44.3 (67)	153	36.6 (71)	67	* *	139	53.2 (53)	153	49.0 (56)
Switzerlan	44	* *	42	* *	17	* *	31	* *	40	* *
Tippecanoe	1,273	23.3 (4)	1,227	18.2 (9)	283	23.7 (33)	1,046	35.1 (17)	1,210	30.7 (21)
Tipton	61	* *	57	* *	17	* *	46	* *	57	* *
Union	48	* *	47	* *	10	* *	35	* *	46	* *
Vanderburg	1,021	39.0 (52)	917	19.3 (13)	347	45.8 (4)	868	22.4 (4)	906	7.9 (3)
Vermillion	89	* *	87	* *	30	* *	72	* *	87	* *
Vigo	676	37.7 (41)	600	23.2 (27)	218	34.4 (13)	560	49.5 (44)	592	42.9 (39)
Wabash	217	37.8 (42)	208	27.4 (49)	79	* *	182	31.9 (13)	206	19.4 (11)
Warren	67	* *	67	* *	29	* *	49	* *	67	* *
Warrick	246	45.1 (70)	236	26.7 (42)	107	31.8 (21)	206	54.4 (56)	233	61.4 (72)
Washington	168	41.1 (59)	163	25.8 (34)	66	* *	145	44.1 (31)	162	27.2 (18)
Wayne	605	38.7 (51)	586	26.5 (40)	222	28.4 (28)	490	51.6 (49)	577	37.8 (29)
Wells	158	41.8 (62)	155	29.0 (58)	65	* *	120	49.2 (43)	153	43.8 (44)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
White	213	31.5 (18)	208	20.7 (16)	66	* *	178	29.8 (10)	207	15.0 (8)
Whitley	184	38.0 (46)	178	21.3 (20)	68	* *	150	44.0 (30)	177	39.0 (33)
Unknown Counties	306	39.2 (53)	3	* *	1	* *	291	38.5 (20)	3	* *
Indiana	46,227	29.0	44,078	20.3	12,540	33.8	35,519	37.4	43,331	31.1
Nation (Prior Year)	814,631	23.0	715,034	12.1	137,883	38.1	660,570	16.3	562,126	15.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams	193	6.7 (19)	5.2 (12)		166	12.0 (26)	146	3.4 (49)	193	82.4 (3)
Allen	2,570	9.1 (54)	4.6 (7)		2,117	15.1 (58)	1,794	3.4 (49)	2,575	64.8 (50)
Bartholome	498	6.4 (14)	6.6 (36)		408	12.3 (32)	358	2.5 (30)	498	70.9 (30)
Benton	39	* * *	* *		31	* *	23	* *	39	* *
Blackford	99	* * *	* *		78	* *	71	* *	99	* *
Boone	189	5.3 (6)	6.9 (43)		151	13.9 (49)	130	2.3 (26)	189	56.1 (67)
Brown	59	* * *	* *		47	* *	39	* *	59	* *
Carroll	79	* * *	* *		65	* *	58	* *	79	* *
Cass	351	5.4 (9)	6.0 (27)		305	9.2 (6)	273	1.5 (13)	355	63.9 (55)
Clark	674	7.6 (36)	5.8 (21)		547	13.9 (49)	469	3.2 (45)	677	62.5 (56)
Clay	225	5.3 (6)	4.9 (8)		192	12.0 (26)	169	1.8 (17)	225	65.3 (46)
Clinton	257	5.4 (9)	8.9 (67)		216	10.6 (13)	192	2.1 (25)	258	72.1 (27)
Crawford	76	* * *	* *		61	* *	58	* *	76	* *
Daviess	177	6.8 (22)	2.8 (3)		149	13.4 (48)	129	3.9 (54)	177	59.3 (64)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dearborn	308	9.4 (60)	9.4 (68)	241	12.9 (39)	208	3.4 (49)	312	64.7 (51)
Decatur	186	11.8 (69)	5.9 (24)	149	17.4 (66)	123	4.1 (57)	186	72.6 (24)
DeKalb	280	7.1 (25)	7.1 (46)	231	12.1 (29)	202	5.0 (66)	281	86.1 (1)
Delaware	802	9.7 (63)	4.4 (6)	641	12.9 (39)	558	4.5 (64)	802	60.3 (62)
Dubois	242	4.5 (4)	5.8 (21)	187	12.8 (35)	163	0.6 (4)	242	76.9 (10)
Elkhart	1,784	7.3 (31)	8.3 (63)	1,556	11.1 (19)	1,377	2.7 (35)	1,795	74.8 (16)
Fayette	179	7.3 (31)	8.4 (65)	137	13.1 (41)	119	0.8 (8)	179	57.5 (66)
Floyd	447	9.4 (60)	5.8 (21)	367	19.3 (68)	293	2.4 (28)	450	65.6 (44)
Fountain	100	12.0 (71)	6.0 (27)	84	* *	69	* *	100	69.0 (38)
Franklin	94	* *	* *	78	* *	72	* *	94	* *
Fulton	129	3.1 (1)	7.8 (58)	107	9.3 (7)	97	* *	129	72.9 (23)
Gibson	182	7.1 (25)	4.9 (8)	141	16.3 (64)	117	1.7 (16)	185	52.4 (69)
Grant	527	10.2 (67)	5.5 (18)	425	13.2 (42)	367	4.9 (65)	529	50.1 (71)
Greene	159	8.8 (52)	6.3 (32)	127	11.0 (17)	113	4.4 (63)	159	64.2 (53)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)	Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Hamilton	649	8.5 (48)	7.1 (46)	509	12.2 (31)	446	3.1 (40)	650	77.5 (7)
Hancock	283	6.4 (14)	7.4 (51)	218	11.9 (25)	192	0.5 (2)	283	76.7 (12)
Harrison	235	6.4 (14)	6.8 (42)	187	16.0 (63)	157	1.9 (19)	235	74.0 (17)
Hendricks	364	6.6 (17)	6.6 (36)	264	9.8 (9)	238	1.3 (11)	364	76.4 (13)
Henry	288	6.9 (24)	7.6 (55)	248	6.5 (1)	230	4.3 (60)	290	70.0 (34)
Howard	657	6.7 (19)	5.2 (12)	546	10.1 (11)	489	2.7 (35)	660	51.5 (70)
Huntington	217	7.8 (38)	5.1 (11)	186	14.5 (51)	158	2.5 (30)	218	67.4 (40)
Jackson	326	7.4 (34)	8.3 (63)	274	15.0 (56)	230	3.0 (39)	328	72.0 (28)
Jasper	206	4.4 (3)	10.2 (70)	160	8.8 (4)	146	0.7 (6)	206	60.7 (61)
Jay	148	10.1 (65)	2.0 (1)	110	11.8 (24)	97	* *	148	64.2 (53)
Jefferson	201	11.4 (68)	7.5 (52)	165	18.8 (67)	130	3.1 (40)	205	72.2 (26)
Jennings	260	8.1 (41)	6.5 (34)	223	14.8 (55)	190	0.5 (2)	260	67.3 (41)
Johnson	693	7.6 (36)	7.6 (55)	509	13.2 (42)	440	2.3 (26)	695	71.1 (29)
Knox	203	8.4 (46)	3.9 (4)	172	12.8 (35)	150	4.0 (55)	204	61.3 (58)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Kosciusko	410	6.6 (17)	6.6 (36)	317	12.3 (32)	278	0.7 (6)	411	73.5 (22)	
LaGrange	130	9.2 (57)	6.9 (43)	104	15.4 (61)	88	* *	130	78.5 (6)	
Lake	3,746	9.1 (54)	5.9 (24)	2,680	14.7 (54)	2,276	3.3 (46)	3,760	69.3 (37)	
LaPorte	806	8.7 (50)	5.7 (19)	644	14.6 (52)	550	3.5 (53)	807	64.9 (48)	
Lawrence	274	9.9 (64)	6.6 (36)	229	9.6 (8)	207	4.3 (60)	274	70.4 (32)	
Madison	906	8.4 (46)	5.4 (16)	706	15.2 (59)	592	1.9 (19)	914	58.8 (65)	
Marion	8,754	8.8 (52)	6.2 (31)	6,142	14.6 (52)	5,224	2.6 (32)	8,805	64.4 (52)	
Marshall	354	4.2 (2)	7.6 (55)	281	8.5 (3)	257	1.9 (19)	354	77.4 (8)	
Martin	55	* *	* *	44	* *	34	* *	55	* *	
Miami	185	8.6 (49)	5.9 (24)	160	15.0 (56)	136	1.5 (13)	185	59.5 (63)	
Monroe	589	8.0 (39)	9.8 (69)	468	11.3 (22)	414	3.1 (40)	590	80.8 (5)	
Montgomery	258	5.8 (11)	11.6 (72)	207	7.2 (2)	191	2.6 (32)	259	69.9 (35)	
Morgan	399	9.3 (59)	8.5 (66)	284	12.0 (26)	250	2.4 (28)	401	73.6 (20)	
Newton	57	* *	* *	40	* *	32	* *	57	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Noble	283	7.1 (25)	5.3 (14)	233	13.3 (45)	202	4.0 (55)	283	73.9 (19)
Ohio	6	* *	* *	4	* *	2	* *	7	* *
Orange	150	5.3 (6)	2.0 (1)	115	10.4 (12)	103	1.9 (19)	150	74.0 (17)
Owen	111	6.3 (13)	7.2 (48)	94	* *	79	* *	111	75.7 (14)
Parke	75	* *	* *	59	* *	51	* *	75	* *
Perry	122	13.9 (72)	5.7 (19)	95	* *	83	* *	122	53.3 (68)
Pike	66	* *	* *	57	* *	41	* *	68	* *
Porter	723	7.3 (31)	8.2 (62)	552	11.2 (20)	488	1.6 (15)	726	73.6 (20)
Posey	93	* *	* *	71	* *	63	* *	93	* *
Pulaski	65	* *	* *	51	* *	44	* *	66	* *
Putnam	210	7.1 (25)	6.7 (41)	180	10.0 (10)	161	3.1 (40)	211	64.9 (48)
Randolph	166	6.0 (12)	6.6 (36)	143	13.3 (45)	124	0.0 (1)	167	61.1 (60)
Ripley	176	9.1 (54)	6.3 (32)	136	13.2 (42)	118	0.8 (8)	176	61.9 (57)
Rush	97	* *	* *	84	* *	72	* *	99	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
Scott	216	8.3 (45)	6.0 (27)	185	16.8 (65)	152	3.3 (46)	218	46.3 (72)
Shelby	304	7.2 (30)	4.9 (8)	241	13.3 (45)	207	4.3 (60)	306	67.6 (39)
Spencer	96	* * *	* *	79	* *	67	* *	96	* *
St Joseph	1,830	8.7 (50)	7.8 (58)	1,490	10.7 (15)	1,326	3.4 (49)	1,837	76.9 (10)
Starke	161	8.1 (41)	7.5 (52)	141	12.8 (35)	123	4.1 (57)	161	70.8 (31)
Steuben	253	7.5 (35)	11.5 (71)	224	11.2 (20)	196	3.1 (40)	256	83.6 (2)
Sullivan	149	8.1 (41)	5.4 (16)	132	10.6 (13)	118	4.2 (59)	149	69.8 (36)
Switzerlan	36	* * *	* *	27	* *	26	* *	37	* *
Tippecanoe	1,170	7.1 (25)	7.5 (52)	966	10.7 (15)	860	3.3 (46)	1,174	75.0 (15)
Tipton	55	* * *	* *	40	* *	36	* *	55	* *
Union	45	* * *	* *	33	* *	30	* *	45	* *
Vanderburg	896	9.2 (57)	6.1 (30)	773	15.3 (60)	651	2.6 (32)	902	66.1 (42)
Vermillion	87	* * *	* *	71	* *	61	* *	87	* *
Vigo	575	8.0 (39)	7.3 (49)	475	12.8 (35)	407	1.2 (10)	583	65.4 (45)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wabash	201	5.0 (5)	6.5 (34)		171	8.8 (4)	155	1.3 (11)	202	70.3 (33)
Warren	65	* *	* *		48	* *	44	* *	65	* *
Warrick	228	9.6 (62)	7.0 (45)		191	15.7 (62)	160	0.6 (4)	229	65.1 (47)
Washington	148	8.1 (41)	8.1 (61)		128	12.5 (34)	112	1.8 (17)	148	72.3 (25)
Wayne	546	6.8 (22)	7.9 (60)		447	11.6 (23)	394	2.0 (24)	547	61.2 (59)
Wells	148	10.1 (65)	4.1 (5)		116	12.1 (29)	102	2.9 (38)	148	77.0 (9)
White	193	6.7 (19)	7.3 (49)		163	11.0 (17)	143	2.8 (37)	196	65.8 (43)
Whitley	169	11.8 (69)	5.3 (14)		139	23.0 (69)	107	1.9 (19)	169	81.1 (4)
Unknown Counties	2	* *	* *		0	* *	0	* *	2	* *
Indiana	42,474	8.2	6.5		33,235	13.2	28,717	2.8	42,656	67.6
Nation (Prior Year)	880,933	8.0	6.7		948,286	10.7	749,107	3.4	871,811	68.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Adams	619	92.4	0.5	6.1	0.0	0.0	0.2	0.8
Allen	8,646	50.9	26.7	14.0	0.1	5.6	2.2	0.6
Bartholome	1,632	80.3	1.8	15.3	0.0	0.6	1.1	0.9
Benton	122	84.4	0.0	13.9	0.0	0.0	0.0	1.6
Blackford	293	96.9	0.3	1.7	0.0	0.0	0.3	0.7
Boone	612	94.0	1.1	2.9	0.0	0.7	0.2	1.1
Brown	209	94.7	0.0	2.4	0.5	1.0	0.0	1.4
Carroll	248	89.1	0.0	10.1	0.0	0.4	0.4	0.0
Cass	1,148	63.6	0.7	32.7	0.0	1.0	0.9	1.1
Clark	2,271	71.1	10.1	14.4	0.2	1.1	2.0	1.1
Clay	703	96.4	0.6	0.7	0.0	0.4	1.1	0.7
Clinton	901	63.6	0.4	34.6	0.3	0.1	0.4	0.4
Crawford	219	96.8	0.0	1.8	0.0	0.0	0.0	1.4
Daviess	555	77.1	0.5	18.0	0.0	0.0	2.2	2.2
Dearborn	957	96.4	0.9	0.3	0.2	0.1	0.6	1.4
Decatur	556	96.6	0.4	1.6	0.2	0.2	0.2	0.9
DeKalb	986	93.9	0.7	3.4	0.2	0.4	0.5	0.8
Delaware	2,621	82.0	12.0	2.5	0.2	0.7	2.2	0.5
Dubois	782	73.0	0.0	23.7	0.1	1.7	0.6	0.9
Elkhart	6,108	54.7	10.0	32.5	0.0	0.9	1.2	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Fayette	649	96.8	1.8	0.2	0.0	0.2	0.6	0.5
Floyd	1,410	76.0	12.5	6.1	0.1	2.2	1.3	1.8
Fountain	356	95.5	0.0	2.2	0.0	0.8	0.3	1.1
Franklin	325	98.5	0.0	0.3	0.0	0.0	0.3	0.9
Fulton	517	89.0	0.8	9.1	0.0	0.0	0.2	1.0
Gibson	578	91.5	3.3	2.4	0.7	0.7	1.0	0.3
Grant	1,756	79.6	10.6	6.5	0.0	0.3	2.1	0.9
Greene	522	97.1	0.4	1.7	0.0	0.4	0.2	0.2
Hamilton	2,047	77.8	6.9	11.0	0.2	2.5	1.3	0.3
Hancock	866	94.3	1.4	1.8	0.1	0.2	0.8	1.3
Harrison	727	93.3	0.3	3.9	0.0	0.4	1.0	1.2
Hendricks	1,273	86.6	4.6	4.1	0.1	3.0	0.1	1.5
Henry	1,003	94.5	0.8	2.1	0.1	0.3	1.2	1.0
Howard	2,164	83.9	9.6	3.9	0.0	0.4	1.6	0.7
Huntington	726	96.3	0.6	1.5	0.0	0.3	0.7	0.7
Jackson	1,070	82.0	1.1	15.6	0.0	0.6	0.0	0.7
Jasper	697	87.8	0.4	10.9	0.0	0.1	0.4	0.3
Jay	466	89.5	0.6	7.9	0.0	0.2	0.0	1.7
Jefferson	733	87.2	1.8	7.8	0.0	0.5	1.5	1.2
Jennings	864	95.0	0.7	2.5	0.6	0.1	0.3	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Johnson	2,227	87.4	1.4	5.4	0.0	3.1	1.1	1.5
Knox	740	93.6	0.7	1.6	0.1	0.3	1.4	2.3
Kosciusko	1,278	75.4	0.8	21.8	0.1	0.4	0.5	1.2
LaGrange	435	82.1	1.1	12.0	0.0	1.1	1.4	2.3
Lake	13,288	31.1	42.6	23.8	0.1	0.5	1.2	0.8
LaPorte	2,735	68.2	18.6	9.9	0.0	0.7	1.9	0.7
Lawrence	931	96.1	0.4	1.4	0.0	1.2	0.3	0.5
Madison	2,920	77.0	12.1	8.3	0.2	0.3	1.4	0.8
Marion	29,030	34.3	39.6	20.6	0.2	3.0	1.5	0.8
Marshall	1,064	74.5	0.5	22.7	0.1	0.5	1.0	0.8
Martin	173	97.7	0.0	1.2	0.0	0.0	0.0	1.2
Miami	670	91.8	1.8	2.2	0.6	0.7	2.8	0.0
Monroe	2,024	81.1	5.7	3.9	0.0	5.9	2.1	1.3
Montgomery	872	87.6	0.8	9.4	0.1	0.3	0.9	0.8
Morgan	1,275	96.9	0.4	1.2	0.0	0.2	0.6	0.8
Newton	185	92.4	0.0	7.0	0.0	0.5	0.0	0.0
Noble	886	75.2	0.3	22.3	0.1	0.8	0.7	0.6
Ohio	29	*	*	*	*	*	*	*
Orange	487	94.7	1.6	1.6	0.0	0.8	0.8	0.4
Owen	388	95.6	0.3	1.8	0.0	1.0	0.5	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Parke	240	96.7	0.0	1.7	0.4	0.0	0.4	0.8
Perry	386	97.2	0.5	0.8	0.0	0.3	0.0	1.3
Pike	226	96.0	0.4	1.8	0.0	0.0	1.3	0.4
Porter	2,326	77.3	7.4	12.2	0.2	1.1	0.3	1.5
Posey	293	93.5	2.4	1.7	0.0	0.3	1.4	0.7
Pulaski	264	95.8	0.0	1.9	0.4	0.8	0.4	0.8
Putnam	732	95.5	0.8	1.5	0.0	0.5	0.8	0.8
Randolph	622	88.3	0.2	7.4	0.2	0.5	1.4	2.1
Ripley	610	93.3	0.0	3.6	0.7	0.0	1.0	1.5
Rush	312	95.8	0.3	1.6	0.6	0.6	0.6	0.3
Scott	720	96.0	0.0	2.2	0.0	0.4	0.7	0.7
Shelby	929	86.5	1.1	11.4	0.0	0.1	0.3	0.5
Spencer	285	90.2	1.1	6.7	0.0	0.0	0.0	2.1
St Joseph	5,934	49.8	28.3	16.5	0.3	1.7	2.4	1.1
Starke	566	94.5	0.2	4.6	0.0	0.2	0.0	0.5
Steuben	840	91.0	0.6	4.9	0.0	1.3	0.4	1.9
Sullivan	442	97.1	0.5	1.4	0.0	0.0	0.2	0.9
Switzerlan	148	94.6	1.4	0.7	0.0	0.0	0.7	2.7
Tippecanoe	3,948	61.5	10.2	19.1	0.4	6.4	1.2	1.1
Tipton	182	91.8	0.5	7.1	0.0	0.0	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Union	143	96.5	0.7	0.7	0.0	0.7	0.0	1.4
Vanderburg	3,081	71.9	18.3	5.0	0.1	1.0	2.1	1.6
Vermillion	303	98.0	0.3	0.0	0.0	0.3	0.3	1.0
Vigo	2,084	86.5	8.8	2.0	0.0	0.9	1.3	0.5
Wabash	634	89.7	0.6	3.6	0.0	0.6	4.1	1.3
Warren	185	97.8	0.0	1.6	0.0	0.0	0.5	0.0
Warrick	777	95.2	1.4	1.0	0.0	0.3	1.3	0.8
Washington	544	96.1	0.2	1.8	0.4	0.4	0.7	0.4
Wayne	1,839	82.7	5.7	6.9	0.3	0.7	3.0	0.8
Wells	461	92.4	0.4	4.3	0.2	0.2	1.3	1.1
White	668	78.7	0.0	19.6	0.1	0.4	0.4	0.6
Whitley	605	97.4	0.3	1.2	0.0	0.3	0.5	0.3
Unknown Counties	308	68.5	17.2	9.1	0.0	1.9	1.0	2.3
Indiana	142,211	63.8	18.2	13.9	0.1	1.8	1.3	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Adams	619	0.5	6.1	13.4	65.9	13.1	0.8	0.2
Allen	8,646	0.2	4.5	12.8	63.2	17.5	1.1	0.7
Bartholome	1,632	0.1	8.1	16.4	59.3	15.0	1.1	0.0
Benton	122	0.0	6.6	11.5	63.9	16.4	1.6	0.0
Blackford	293	0.0	5.1	18.1	64.5	11.9	0.3	0.0
Boone	612	0.2	4.2	14.4	65.7	14.2	1.3	0.0
Brown	209	0.0	6.2	15.8	59.8	15.3	2.9	0.0
Carroll	248	0.4	4.0	11.3	61.3	22.6	0.4	0.0
Cass	1,148	0.9	5.1	13.4	60.5	18.9	1.0	0.2
Clark	2,271	0.2	5.5	15.7	61.5	16.2	0.7	0.1
Clay	703	0.1	8.0	15.4	63.7	11.9	0.9	0.0
Clinton	901	0.0	6.5	13.2	63.2	15.8	1.3	0.0
Crawford	219	0.0	5.5	17.8	64.8	11.9	0.0	0.0
Daviess	555	0.0	5.0	13.3	65.9	13.9	0.7	1.1
Dearborn	957	0.5	4.5	15.0	64.3	14.3	1.0	0.3
Decatur	556	0.0	8.3	15.8	60.4	14.7	0.7	0.0
DeKalb	986	0.3	5.2	16.4	64.3	12.8	0.9	0.1
Delaware	2,621	0.2	5.5	14.4	64.6	14.2	0.9	0.1
Dubois	782	0.3	4.2	11.6	62.4	19.8	1.5	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Elkhart	6,108	0.4	5.8	13.9	58.6	19.8	1.3	0.0
Fayette	649	0.3	9.6	16.6	60.9	11.9	0.6	0.2
Floyd	1,410	0.4	6.3	16.2	61.8	14.6	0.4	0.3
Fountain	356	0.0	5.3	19.7	59.3	15.7	0.0	0.0
Franklin	325	0.0	6.8	14.5	64.0	13.5	1.2	0.0
Fulton	517	0.2	4.8	19.3	60.3	14.3	1.0	0.0
Gibson	578	0.0	7.1	14.5	65.1	12.5	0.7	0.2
Grant	1,756	0.7	7.5	15.9	63.9	11.4	0.6	0.1
Greene	522	0.2	5.4	18.0	64.2	11.9	0.4	0.0
Hamilton	2,047	0.1	3.1	12.1	59.0	23.6	1.5	0.6
Hancock	866	0.1	5.4	15.5	62.1	16.1	0.8	0.0
Harrison	727	0.0	5.0	14.0	66.2	14.3	0.3	0.3
Hendricks	1,273	0.1	5.2	14.5	59.8	19.1	1.3	0.1
Henry	1,003	0.6	6.0	15.5	64.2	13.2	0.5	0.1
Howard	2,164	0.3	5.6	15.7	63.2	14.1	0.8	0.3
Huntington	726	0.1	3.9	19.1	64.7	10.3	0.4	1.4
Jackson	1,070	0.1	8.4	14.9	63.0	12.7	0.7	0.3
Jasper	697	0.0	5.7	15.2	60.1	17.1	1.7	0.1
Jay	466	0.0	6.2	14.4	64.2	14.8	0.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Jefferson	733	0.5	6.7	16.1	61.8	14.1	0.8	0.0
Jennings	864	0.2	8.9	21.1	56.4	13.1	0.3	0.0
Johnson	2,227	0.1	5.9	14.8	61.9	16.1	1.2	0.0
Knox	740	0.3	6.5	16.2	62.0	14.1	0.4	0.5
Kosciusko	1,278	0.3	6.7	15.2	58.2	18.4	1.3	0.0
LaGrange	435	0.2	6.0	13.6	58.6	19.1	1.8	0.7
Lake	13,288	0.4	5.9	12.8	58.3	21.1	1.4	0.1
LaPorte	2,735	0.3	5.5	13.5	60.6	18.9	1.2	0.0
Lawrence	931	0.2	9.0	17.0	61.0	11.6	1.2	0.0
Madison	2,920	0.4	5.9	15.0	63.6	14.1	0.9	0.1
Marion	29,030	0.3	5.2	12.2	61.6	19.5	1.2	0.1
Marshall	1,064	0.1	5.3	12.1	57.7	22.7	1.9	0.3
Martin	173	0.0	9.8	13.9	61.8	13.9	0.6	0.0
Miami	670	0.4	6.0	17.8	60.7	14.5	0.4	0.1
Monroe	2,024	0.1	3.3	13.7	63.2	17.8	1.4	0.6
Montgomery	872	0.3	6.7	15.3	60.2	16.1	1.5	0.0
Morgan	1,275	0.2	5.8	14.6	64.5	13.9	0.9	0.2
Newton	185	0.5	3.2	13.0	65.9	16.8	0.5	0.0
Noble	886	0.2	5.2	14.2	61.9	17.6	0.8	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Ohio	29	*	*	*	*	*	*	*
Orange	487	0.6	4.7	15.6	61.6	17.0	0.4	0.0
Owen	388	0.3	5.4	17.3	63.4	12.1	1.5	0.0
Parke	240	0.0	8.8	18.8	58.3	13.3	0.8	0.0
Perry	386	0.5	8.0	16.6	61.4	13.0	0.5	0.0
Pike	226	0.0	6.2	17.3	61.9	14.2	0.4	0.0
Porter	2,326	0.0	4.5	12.0	62.5	19.3	1.5	0.2
Posey	293	0.0	6.5	16.7	62.1	14.0	0.7	0.0
Pulaski	264	0.4	4.2	17.0	64.4	12.9	0.8	0.4
Putnam	732	0.5	6.6	15.7	65.0	11.5	0.7	0.0
Randolph	622	0.3	5.1	14.8	63.0	15.8	0.6	0.3
Ripley	610	0.0	6.4	16.1	61.5	15.2	0.8	0.0
Rush	312	0.6	7.4	16.3	64.7	10.3	0.6	0.0
Scott	720	0.4	6.3	16.8	66.3	9.3	1.0	0.0
Shelby	929	0.2	6.4	15.5	63.0	13.8	1.2	0.0
Spencer	285	0.4	5.6	14.7	61.8	16.5	1.1	0.0
St Joseph	5,934	0.2	6.3	13.6	59.1	19.5	1.3	0.0
Starke	566	0.9	5.3	14.3	61.5	16.6	1.2	0.2
Steuben	840	0.1	5.1	16.2	64.4	13.5	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sullivan	442	0.7	8.4	18.8	57.7	13.8	0.7	0.0
Switzerland	148	0.0	7.4	14.2	60.8	14.9	2.7	0.0
Tippecanoe	3,948	0.2	5.6	13.1	59.7	20.0	1.2	0.2
Tipton	182	0.0	11.0	13.7	61.0	13.2	1.1	0.0
Union	143	0.0	5.6	16.8	58.0	18.2	1.4	0.0
Vanderburg	3,081	0.4	6.3	14.0	64.6	14.3	0.5	0.0
Vermillion	303	0.0	5.0	18.5	64.4	11.6	0.7	0.0
Vigo	2,084	0.4	8.1	16.2	62.3	12.1	0.8	0.0
Wabash	634	0.2	4.6	15.0	67.2	11.5	1.3	0.3
Warren	185	0.0	6.5	18.9	60.0	14.1	0.5	0.0
Warrick	777	0.0	4.5	15.2	64.7	15.3	0.0	0.3
Washington	544	0.0	7.2	18.6	62.1	10.8	0.9	0.4
Wayne	1,839	0.2	5.5	14.6	64.2	14.9	0.7	0.0
Wells	461	0.7	4.1	14.5	65.5	14.5	0.7	0.0
White	668	0.7	6.9	13.5	60.6	16.3	1.9	0.0
Whitley	605	0.3	4.3	15.4	65.8	12.9	1.3	0.0
Unknown Counties	308	0.3	6.5	11.7	65.6	14.6	1.3	0.0
Indiana	142,211	0.3	5.7	14.0	61.6	17.3	1.1	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams	497	7.0 (79)	53.3 (49)	474	16.5 (14)	55.1 (57)	82	* *	129	27.9(41)	599	0.7 (10)	599	2.0(19)
Allen	7,044	4.7 (28)	53.9 (54)	6,715	20.7 (81)	50.7 (15)	1,254	41.9 (36)	1,305	31.6(51)	8,380	2.5 (39)	8,380	3.8(48)
Bartholome	1,291	7.5 (86)	52.9 (41)	1,240	16.9 (19)	55.5 (62)	259	32.4 (15)	250	26.4(32)	1,590	5.3 (81)	1,590	6.6(78)
Benton	96	* *	* *	92	* *	* *	15	* *	26	* *	118	7.6 (89)	118	5.1(60)
Blackford	237	6.3 (68)	49.4 (10)	231	19.5 (69)	48.9 (6)	56	* *	46	* *	289	3.1 (49)	289	9.0(87)
Boone	464	6.7 (74)	51.7 (28)	438	17.4 (26)	51.6 (23)	85	* *	155	21.3(15)	586	1.0 (13)	586	1.2(11)
Brown	167	6.0 (61)	43.7 (1)	159	18.9 (53)	50.9 (16)	32	* *	56	* *	202	7.9 (90)	202	5.9(70)
Carroll	199	3.0 (4)	56.3 (76)	193	17.6 (29)	48.2 (4)	34	* *	43	* *	246	0.8 (11)	246	2.0(19)
Cass	917	4.6 (23)	55.0 (62)	845	20.6 (78)	48.2 (4)	171	29.8 (11)	135	20.0(10)	1,082	3.3 (55)	1,082	1.9(15)
Clark	1,784	4.6 (23)	56.8 (79)	1,693	19.2 (61)	52.4 (33)	367	34.6 (21)	425	31.3(49)	2,188	6.5 (88)	2,188	2.5(36)
Clay	589	4.9 (36)	58.6 (86)	561	20.1 (75)	53.1 (36)	82	* *	123	18.7 (8)	684	2.8 (46)	684	4.1(50)
Clinton	753	3.9 (10)	62.8 (91)	724	19.3 (62)	49.9 (10)	171	33.9 (18)	155	23.2(23)	882	2.3 (34)	882	2.6(41)
Crawford	175	5.1 (38)	57.1 (80)	166	20.5 (77)	51.8 (26)	38	* *	34	* *	212	1.9 (28)	212	2.4(34)
Daviess	444	2.9 (3)	61.0 (88)	400	20.8 (82)	50.5 (14)	92	* *	315	37.8(58)	514	2.5 (39)	514	8.0(82)
Dearborn	756	5.8 (57)	48.9 (6)	719	14.5 (4)	57.6 (80)	146	48.6 (40)	210	23.3(24)	922	1.5 (19)	922	2.2(25)
Decatur	434	6.7 (74)	55.8 (70)	419	17.4 (26)	51.3 (20)	53	* *	124	14.5 (2)	540	3.9 (67)	540	5.9(70)
DeKalb	788	5.2 (40)	50.6 (16)	746	19.3 (62)	54.6 (51)	160	28.8 (6)	130	23.8(25)	953	0.9 (12)	953	2.3(33)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Delaware	1,976	5.4 (50)	54.4 (59)	1,860	18.9 (53)	56.2 (72)	388	40.7 (33)	311	31.2(48)	2,515	2.1 (33)	2,515	2.0(19)
Dubois	589	5.1 (38)	61.1 (89)	570	17.7 (32)	53.5 (42)	135	26.7 (3)	133	17.3 (5)	767	5.0 (78)	767	3.1(44)
Elkhart	5,123	3.9 (10)	57.4 (81)	4,883	19.7 (73)	51.8 (26)	623	27.1 (4)	969	20.6(13)	5,897	3.1 (49)	5,897	3.1(44)
Fayette	525	7.2 (81)	52.0 (30)	504	17.3 (23)	58.1 (83)	87	* *	105	33.3(55)	631	1.1 (15)	631	2.1(23)
Floyd	1,093	5.3 (45)	53.8 (52)	1,019	19.4 (66)	52.3 (30)	201	34.8 (22)	409	25.4(30)	1,342	8.3 (91)	1,342	12.4(91)
Fountain	287	2.4 (1)	55.1 (63)	273	19.0 (55)	60.4 (88)	52	* *	63	* *	343	5.2 (80)	343	5.5(68)
Franklin	249	5.2 (40)	57.4 (81)	237	19.0 (55)	55.7 (67)	40	* *	59	* *	316	1.9 (28)	316	2.2(25)
Fulton	423	4.3 (16)	52.2 (35)	390	18.5 (44)	58.2 (84)	71	* *	113	28.3(42)	491	0.0 (1)	491	0.2(2)
Gibson	437	5.3 (45)	53.1 (45)	407	18.7 (50)	55.5 (62)	81	* *	101	22.8(20)	549	1.6 (21)	549	2.0(19)
Grant	1,419	5.9 (59)	51.4 (26)	1,337	19.3 (62)	53.3 (39)	279	45.9 (39)	487	26.9(36)	1,684	2.7 (44)	1,684	8.3(84)
Greene	411	6.1 (65)	52.1 (31)	390	20.8 (82)	55.6 (64)	82	* *	121	24.0(27)	509	3.1 (49)	509	1.0(10)
Hamilton	1,539	4.5 (21)	51.3 (24)	1,465	19.1 (58)	54.1 (46)	384	40.6 (32)	549	31.5(50)	1,984	3.9 (67)	1,984	4.1(50)
Hancock	651	5.7 (54)	52.1 (31)	614	14.7 (5)	59.1 (87)	149	28.9 (8)	193	16.1 (3)	834	3.5 (57)	834	2.4(34)
Harrison	565	5.0 (37)	52.9 (41)	528	21.8 (85)	50.0 (11)	78	* *	101	17.8 (6)	694	2.7 (44)	694	2.6(41)
Hendricks	942	4.7 (28)	52.1 (31)	801	12.2 (2)	57.7 (82)	248	41.5 (34)	674	39.6(61)	1,136	2.4 (36)	1,136	2.5(36)
Henry	828	6.8 (77)	50.7 (18)	797	18.2 (40)	57.5 (79)	127	31.5 (14)	127	18.9 (9)	979	1.8 (27)	979	0.9(9)
Howard	1,761	5.2 (40)	51.8 (29)	1,699	19.4 (66)	50.9 (16)	309	41.7 (35)	373	32.2(53)	2,106	1.6 (21)	2,106	2.2(25)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Huntington	624	8.3 (88)	49.7 (12)	596	19.5 (69)	54.5 (49)	75	* *	91	* *	702	3.8 (66)	702	9.7(89)
Jackson	896	6.0 (61)	54.9 (61)	862	19.1 (58)	52.1 (28)	167	38.3 (27)	218	23.9(26)	1,047	3.2 (54)	1,047	5.3(64)
Jasper	540	4.8 (32)	50.0 (13)	492	16.5 (14)	53.7 (45)	60	* *	95	* *	652	0.3 (5)	652	0.5(5)
Jay	363	5.8 (57)	55.1 (63)	345	19.7 (73)	53.3 (39)	69	* *	81	* *	450	3.1 (49)	450	2.2(25)
Jefferson	581	5.7 (54)	54.2 (56)	547	16.3 (12)	52.3 (30)	100	31.0 (12)	126	11.1 (1)	704	5.4 (83)	704	6.3(74)
Jennings	756	6.7 (74)	53.0 (44)	727	19.5 (69)	51.6 (23)	77	* *	118	28.8(43)	840	1.0 (13)	840	1.7(14)
Johnson	1,628	6.1 (65)	51.1 (23)	1,546	16.5 (14)	56.0 (68)	377	35.5 (23)	424	20.8(14)	2,156	3.5 (57)	2,156	7.9(81)
Knox	606	7.3 (84)	53.3 (49)	517	17.0 (20)	56.1 (69)	99	* *	380	21.6(18)	654	3.7 (62)	654	4.1(50)
Kosciusko	1,001	4.7 (28)	55.5 (68)	926	17.8 (35)	53.1 (36)	245	33.9 (18)	336	21.4(16)	1,205	4.5 (74)	1,205	5.0(59)
LaGrange	337	5.3 (45)	50.1 (14)	316	18.4 (41)	55.1 (57)	74	* *	122	23.0(21)	412	4.1 (72)	412	6.1(72)
Lake	9,445	4.2 (13)	57.6 (83)	8,430	18.5 (44)	54.9 (56)	2,055	44.9 (37)	1,597	30.7(45)	12,341	2.4 (36)	12,341	4.3(54)
LaPorte	2,146	5.2 (40)	54.5 (60)	1,962	16.1 (9)	54.8 (54)	444	34.0 (20)	793	26.4(32)	2,567	2.5 (39)	2,567	2.1(23)
Lawrence	746	6.4 (70)	53.6 (51)	704	17.6 (29)	51.4 (21)	103	49.5 (41)	176	32.4(54)	901	4.9 (77)	901	7.7(80)
Madison	2,169	4.7 (28)	55.4 (66)	2,076	17.3 (23)	53.1 (36)	529	45.6 (38)	587	34.1(56)	2,843	1.9 (28)	2,843	2.5(36)
Marion	20,240	4.4 (19)	56.7 (78)	18,550	19.1 (58)	53.3 (39)	4,409	38.1 (26)	4,342	27.6(40)	27,472	3.9 (67)	27,472	5.3(64)
Marshall	856	3.2 (6)	61.1 (89)	814	16.3 (12)	56.3 (73)	187	23.0 (2)	165	17.0 (4)	1,029	5.7 (85)	1,029	6.5(77)
Martin	143	4.2 (13)	51.0 (22)	137	11.7 (1)	62.0 (89)	21	* *	14	* *	166	1.2 (17)	166	5.4(67)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	556	6.5 (71)	52.3 (38)	520	17.7 (32)	55.2 (59)	95	* *	401	25.9(31)	637	2.8 (46)	637	5.2(62)
Monroe	1,628	5.6 (53)	49.4 (10)	1,520	17.9 (36)	52.3 (30)	297	29.3 (10)	464	24.6(28)	1,919	2.4 (36)	1,919	1.9(15)
Montgomery	717	6.1 (65)	52.6 (39)	681	17.9 (36)	57.6 (80)	140	28.6 (5)	194	21.6(18)	843	3.7 (62)	843	4.7(57)
Morgan	926	6.6 (73)	55.3 (65)	872	17.3 (23)	55.2 (59)	173	40.5 (31)	224	30.8(46)	1,228	1.7 (24)	1,228	2.2(25)
Newton	141	5.7 (54)	48.2 (4)	136	17.6 (29)	47.1 (1)	23	* *	26	* *	180	0.0 (1)	180	0.6(6)
Noble	701	4.6 (23)	53.1 (45)	660	17.7 (32)	53.5 (42)	158	39.2 (28)	174	31.0(47)	846	0.5 (8)	846	1.5(13)
Ohio	22	* *	* *	22	* *	* *	5	* *	2	* *	29	* *	29	* *
Orange	411	4.4 (19)	60.6 (87)	393	22.6 (86)	48.1 (3)	52	* *	56	* *	471	4.5 (74)	471	11.7(90)
Owen	334	7.2 (81)	52.1 (31)	323	17.0 (20)	51.7 (25)	51	* *	78	* *	377	6.4 (87)	377	6.4(75)
Parke	199	6.5 (71)	49.2 (9)	197	16.2 (10)	56.9 (76)	20	* *	37	* *	240	1.7 (24)	240	2.5(36)
Perry	289	5.2 (40)	46.7 (2)	276	19.6 (72)	50.0 (11)	59	* *	72	* *	374	3.5 (57)	374	5.3(64)
Pike	175	4.6 (23)	54.3 (57)	165	18.8 (51)	58.2 (84)	31	* *	26	* *	216	3.7 (62)	216	8.3(84)
Porter	1,744	4.1 (12)	52.2 (35)	1,640	17.5 (28)	54.6 (51)	351	28.8 (6)	419	25.1(29)	2,233	3.3 (55)	2,233	3.0(43)
Posey	226	5.3 (45)	55.8 (70)	207	15.9 (8)	55.6 (64)	52	* *	57	* *	275	0.4 (6)	275	0.4(4)
Pulaski	199	3.0 (4)	50.8 (20)	178	18.5 (44)	55.6 (64)	24	* *	27	* *	243	0.4 (6)	243	0.0(1)
Putnam	605	4.3 (16)	53.9 (54)	572	16.8 (18)	58.6 (86)	80	* *	158	29.1(44)	706	5.8 (86)	706	8.8(86)
Randolph	501	9.0 (91)	53.1 (45)	478	18.6 (48)	54.2 (47)	115	37.4 (25)	101	26.7(34)	605	1.2 (17)	605	4.5(56)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight	%	< Ideal	> Ideal	3rd Trimester		Postpartum						
						Number	% (Rank)	%	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ripley	467	4.3 (16)	56.5 (77)	437	18.1 (38)	54.7 (53)	72	* *	119	38.7 (60)	582	5.0 (78)	582	9.5 (88)
Rush	253	7.1 (80)	49.0 (7)	244	18.4 (41)	54.5 (49)	37	* *	72	* *	305	3.9 (67)	305	5.2 (62)
Scott	597	7.4 (85)	53.1 (45)	573	23.9 (88)	51.1 (19)	75	* *	123	26.8 (35)	697	3.6 (60)	697	4.7 (57)
Shelby	703	6.0 (61)	54.3 (57)	665	17.1 (22)	56.1 (69)	134	40.3 (30)	423	27.0 (37)	894	2.0 (31)	894	1.9 (15)
Spencer	223	7.6 (87)	53.8 (52)	215	18.1 (38)	49.3 (8)	53	* *	30	* *	281	2.5 (39)	281	3.2 (46)
St Joseph	4,704	4.6 (23)	55.5 (68)	4,448	18.5 (44)	53.5 (42)	906	39.3 (29)	1,284	31.8 (52)	5,710	1.5 (19)	5,710	2.2 (25)
Starke	466	5.4 (50)	50.9 (21)	446	15.7 (7)	56.1 (69)	60	* *	64	* *	550	2.5 (39)	550	4.4 (55)
Steuben	704	4.8 (32)	55.8 (70)	677	21.4 (84)	52.6 (34)	76	* *	108	20.4 (11)	816	1.1 (15)	816	0.6 (6)
Sullivan	382	3.7 (7)	57.6 (83)	356	18.8 (51)	54.8 (54)	61	* *	164	37.8 (58)	419	0.2 (4)	419	1.9 (15)
Switzerlan	117	6.0 (61)	51.3 (24)	111	12.6 (3)	64.0 (90)	17	* *	33	* *	141	0.0 (1)	141	0.7 (8)
Tiptecanoe	3,240	4.8 (32)	50.7 (18)	3,060	19.3 (62)	51.5 (22)	512	32.8 (16)	466	20.4 (11)	3,805	2.0 (31)	3,805	1.4 (12)
Tipton	127	8.7 (89)	55.9 (74)	119	24.4 (90)	47.1 (1)	35	* *	41	* *	175	1.7 (24)	175	8.0 (82)
Union	107	2.8 (2)	57.9 (85)	100	24.0 (89)	49.0 (7)	21	* *	36	* *	136	3.7 (62)	136	2.2 (25)
Vanderburg	2,576	4.5 (21)	55.4 (66)	2,174	18.4 (41)	56.8 (75)	513	37.2 (24)	1,801	23.0 (21)	2,690	2.8 (46)	2,690	4.2 (53)
Vermillion	249	7.2 (81)	49.0 (7)	240	16.7 (17)	56.3 (73)	61	* *	43	* *	297	4.4 (73)	297	6.1 (72)
Vigo	1,743	5.3 (45)	52.8 (40)	1,467	16.2 (10)	57.0 (77)	312	33.0 (17)	1,050	27.0 (37)	1,819	2.3 (34)	1,819	2.5 (36)
Wabash	510	6.9 (78)	52.2 (35)	469	18.6 (48)	51.0 (18)	100	18.0 (1)	407	17.9 (7)	595	4.5 (74)	595	6.4 (75)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
Warren	144	4.2 (13)	47.9 (3)	139	19.4 (66)	55.4 (61)	20	* *	57	* *	180	5.6 (84)	180	2.2(25)
Warrick	605	6.3 (68)	52.9 (41)	569	14.9 (6)	57.3 (78)	120	29.2 (9)	136	27.2(39)	744	3.1 (49)	744	5.5(68)
Washington	463	3.7 (7)	56.2 (75)	446	20.6 (78)	49.3 (8)	52	* *	56	* *	530	4.0 (71)	530	7.0(79)
Wayne	1,521	5.5 (52)	55.8 (70)	1,443	22.6 (86)	50.4 (13)	224	31.3 (13)	296	34.8(57)	1,783	5.3 (81)	1,783	4.0(49)
Wells	353	3.7 (7)	51.6 (27)	336	19.0 (55)	54.2 (47)	75	* *	74	* *	449	1.6 (21)	449	5.1(60)
White	546	4.8 (32)	50.5 (15)	529	20.6 (78)	52.2 (29)	74	* *	89	* *	650	0.6 (9)	650	0.3(3)
Whitley	476	5.9 (59)	50.6 (16)	455	20.2 (76)	52.7 (35)	89	* *	131	21.4(16)	585	3.6 (60)	585	3.2(46)
Unknown Counties	295	8.8 (90)	48.5 (5)	1	* *	* *	78	* *	3	* *	4	* *	4	* *
Indiana	109,475	5.0	54.7	101,695	18.7	53.5	21,025	37.0	27,836	26.9	135,164	3.0	135,164	4.1

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams	409	42.1 (83)	55.5 (83)	499	52.5 (15)	498	24.1 (4)	264	91.3 (13)	
Allen	5,303	57.8 (63)	35.8 (61)	7,063	44.7 (37)	7,043	14.8 (56)	3,505	89.5 (32)	
Bartholome	1,066	56.0 (68)	35.4 (59)	1,299	36.0 (79)	1,291	18.4 (23)	661	90.3 (22)	
Benton	77	* *	* *	93	* *	96	* *	52	* *	
Blackford	185	65.9 (40)	26.5 (33)	240	42.9 (50)	239	18.4 (23)	116	90.5 (20)	
Boone	364	63.2 (47)	31.3 (47)	468	44.4 (40)	465	11.8 (83)	227	91.2 (15)	
Brown	130	56.2 (66)	39.2 (67)	168	42.3 (53)	168	28.6 (2)	79	* *	
Carroll	144	67.4 (34)	25.7 (29)	201	41.8 (54)	199	16.1 (43)	95	* *	
Cass	685	59.7 (59)	33.9 (56)	918	43.8 (45)	919	13.2 (73)	478	78.5 (75)	
Clark	1,358	62.2 (48)	31.9 (49)	1,793	40.2 (59)	1,786	17.9 (28)	890	91.3 (13)	
Clay	404	57.7 (64)	38.1 (66)	592	54.9 (6)	589	17.3 (31)	298	86.6 (52)	
Clinton	579	58.7 (61)	32.3 (50)	750	37.2 (72)	754	18.8 (18)	359	89.4 (34)	
Crawford	122	61.5 (51)	27.0 (35)	176	48.9 (25)	175	15.4 (50)	92	* *	
Daviess	276	78.3 (5)	14.9 (3)	447	41.4 (56)	443	12.6 (76)	241	80.9 (72)	
Dearborn	476	71.8 (18)	23.1 (21)	764	40.2 (59)	755	15.8 (47)	391	90.3 (22)	
Decatur	342	60.8 (54)	35.4 (59)	443	46.3 (29)	432	14.1 (64)	224	81.3 (71)	
DeKalb	560	67.9 (30)	28.6 (38)	787	43.2 (47)	790	13.9 (65)	393	90.6 (18)	
Delaware	1,389	72.4 (14)	22.6 (18)	1,993	43.8 (45)	1,969	17.9 (28)	1,019	90.5 (20)	
Dubois	399	78.4 (4)	17.0 (6)	596	34.2 (85)	589	21.6 (10)	293	86.3 (54)	
Elkhart	4,372	42.2 (82)	53.4 (82)	5,125	53.5 (12)	5,135	19.8 (15)	2,567	81.5 (70)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Fayette	320	73.8 (11)	21.3 (12)	521	49.5 (22)	525	12.2 (81)	237	84.0 (66)	
Floyd	787	60.5 (58)	33.2 (54)	1,116	39.3 (64)	1,096	19.5 (17)	568	89.1 (37)	
Fountain	234	47.0 (78)	46.6 (77)	290	45.2 (34)	286	16.1 (43)	141	85.1 (60)	
Franklin	168	70.8 (22)	26.2 (31)	252	40.5 (58)	249	13.3 (71)	126	89.7 (30)	
Fulton	293	67.6 (32)	27.6 (36)	426	43.0 (49)	423	14.9 (55)	210	89.0 (39)	
Gibson	296	75.0 (9)	22.0 (15)	449	44.3 (41)	439	14.6 (57)	234	93.6 (4)	
Grant	930	73.1 (12)	20.5 (10)	1,420	39.8 (63)	1,422	12.8 (75)	693	90.0 (25)	
Greene	238	72.7 (13)	20.2 (8)	418	44.7 (37)	320	16.9 (36)	190	90.0 (25)	
Hamilton	1,075	71.8 (18)	22.7 (19)	1,556	34.6 (83)	1,541	20.2 (14)	803	91.9 (9)	
Hancock	460	65.4 (43)	29.8 (44)	654	36.7 (74)	653	19.8 (15)	349	91.7 (11)	
Harrison	371	56.1 (67)	36.9 (65)	567	50.3 (20)	566	20.3 (13)	288	95.1 (1)	
Hendricks	596	78.7 (3)	15.1 (5)	953	30.3 (89)	935	22.4 (8)	498	92.6 (5)	
Henry	642	67.0 (35)	29.8 (44)	835	49.3 (23)	832	13.9 (65)	402	81.6 (69)	
Howard	1,347	60.7 (55)	32.4 (52)	1,769	43.1 (48)	1,762	10.5 (87)	892	86.4 (53)	
Huntington	477	40.5 (84)	56.4 (84)	624	63.9 (2)	624	18.3 (26)	303	84.2 (63)	
Jackson	670	70.0 (24)	23.7 (22)	905	44.2 (43)	897	14.6 (57)	457	92.1 (8)	
Jasper	390	51.3 (72)	43.6 (73)	534	53.7 (10)	535	16.4 (38)	259	84.6 (61)	
Jay	266	72.2 (15)	23.7 (22)	368	49.2 (24)	364	14.3 (63)	176	90.3 (22)	
Jefferson	480	60.6 (56)	36.3 (63)	589	46.9 (27)	583	14.4 (62)	288	86.1 (55)	
Jennings	624	28.0 (85)	68.3 (85)	756	65.7 (1)	756	12.3 (79)	362	77.9 (76)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	1,303	66.2 (39)	26.3 (32)		1,636	31.7 (88)	1,610	17.2 (33)	795	89.6 (31)
Knox	368	76.9 (6)	20.4 (9)		617	44.2 (43)	610	7.2 (89)	336	89.3 (35)
Kosciusko	732	76.8 (8)	14.9 (3)		1,005	29.3 (90)	1,002	16.4 (38)	542	88.0 (42)
LaGrange	227	79.7 (2)	12.8 (2)		342	36.0 (79)	337	16.3 (40)	182	87.4 (46)
Lake	6,341	69.6 (26)	21.5 (13)		9,516	34.4 (84)	8,941	15.5 (49)	4,657	87.4 (46)
LaPorte	1,618	58.7 (61)	33.1 (53)		2,159	39.0 (65)	1,889	20.5 (11)	1,049	93.7 (3)
Lawrence	555	56.4 (65)	39.3 (68)		749	53.8 (9)	744	13.4 (69)	360	82.2 (68)
Madison	1,698	61.1 (53)	24.2 (25)		2,173	36.4 (77)	2,172	14.6 (57)	1,127	89.5 (32)
Marion	14,872	69.3 (27)	21.9 (14)		20,267	32.4 (87)	20,012	11.6 (84)	10,402	86.9 (50)
Marshall	688	65.7 (42)	28.6 (38)		860	36.5 (75)	857	18.4 (23)	444	87.6 (44)
Martin	94	* *	* *		142	54.9 (6)	143	21.7 (9)	72	* *
Miami	370	68.4 (28)	29.2 (40)		561	48.1 (26)	556	18.0 (27)	273	87.9 (43)
Monroe	1,142	63.3 (46)	32.3 (50)		1,635	46.1 (31)	1,088	29.1 (1)	779	91.4 (12)
Montgomery	621	46.1 (80)	48.0 (78)		719	44.9 (35)	719	15.9 (45)	344	89.2 (36)
Morgan	691	64.7 (44)	29.5 (42)		933	36.4 (77)	927	18.8 (18)	461	90.0 (25)
Newton	108	67.6 (32)	24.1 (24)		138	39.9 (62)	141	11.3 (86)	72	* *
Noble	497	72.2 (15)	20.7 (11)		705	38.0 (68)	700	13.4 (69)	376	89.1 (37)
Ohio	14	* *	* *		22	* *	22	* *	9	* *
Orange	310	52.9 (70)	45.5 (74)		409	57.0 (5)	410	17.8 (30)	205	84.4 (62)
Owen	242	51.2 (73)	46.3 (76)		333	51.7 (16)	331	13.9 (65)	156	76.9 (77)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Parke	152	52.6 (71)	42.8 (72)		198	50.5 (19)	178	23.0 (7)	89	* *
Perry	238	62.2 (48)	33.2 (54)		294	42.5 (51)	291	2.4 (91)	160	94.4 (2)
Pike	99	* *	* *		177	41.8 (54)	175	12.6 (76)	93	* *
Porter	1,177	68.3 (29)	26.7 (34)		1,753	40.6 (57)	1,742	23.7 (5)	908	90.0 (25)
Posey	162	66.7 (36)	29.6 (43)		229	45.4 (33)	226	15.9 (45)	118	87.3 (48)
Pulaski	129	65.9 (40)	27.9 (37)		198	51.5 (17)	198	18.7 (21)	85	* *
Putnam	463	46.4 (79)	50.1 (80)		604	54.0 (8)	606	17.0 (34)	291	85.6 (58)
Randolph	323	83.3 (1)	11.1 (1)		506	35.8 (82)	500	15.8 (47)	256	92.2 (6)
Ripley	344	63.7 (45)	31.4 (48)		468	40.0 (61)	470	13.0 (74)	239	84.1 (65)
Rush	165	69.7 (25)	26.1 (30)		252	51.2 (18)	252	9.1 (88)	127	75.6 (78)
Scott	495	59.2 (60)	36.6 (64)		596	53.7 (10)	598	15.1 (53)	296	83.1 (67)
Shelby	492	67.9 (30)	24.8 (26)		705	36.5 (75)	709	12.1 (82)	380	87.1 (49)
Spencer	171	66.7 (36)	29.2 (40)		220	35.9 (81)	223	7.2 (89)	126	85.7 (57)
St Joseph	3,568	71.3 (20)	22.5 (17)		4,742	37.5 (70)	4,708	16.2 (42)	2,409	91.2 (15)
Starke	370	53.5 (69)	41.9 (70)		463	52.7 (14)	467	15.4 (50)	230	80.9 (72)
Steuben	517	47.2 (77)	49.9 (79)		709	62.2 (3)	701	12.6 (76)	366	80.6 (74)
Sullivan	196	74.5 (10)	22.4 (16)		381	46.2 (30)	381	16.3 (40)	203	85.2 (59)
Switzerlan	90	* *	* *		118	39.0 (65)	117	15.4 (50)	51	* *
Tippecanoe	2,690	47.6 (76)	45.7 (75)		3,238	50.1 (21)	3,009	23.5 (6)	1,574	86.1 (55)
Tipton	91	* *	* *		129	38.0 (68)	128	18.8 (18)	78	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Union	72	* *	* *	* *	109	38.5 (67)	107	18.7 (21)	50	* *
Vanderburg	1,699	71.1 (21)	22.7 (19)		2,621	37.5 (70)	2,586	11.6 (84)	1,314	91.2 (15)
Vermillion	168	72.0 (17)	25.6 (28)		253	37.2 (72)	249	20.5 (11)	123	91.9 (9)
Vigo	953	76.9 (6)	17.9 (7)		1,758	42.5 (51)	1,746	12.3 (79)	908	88.2 (41)
Wabash	360	66.7 (36)	30.6 (46)		517	44.3 (41)	512	14.6 (57)	266	90.6 (18)
Warren	123	49.6 (75)	41.5 (69)		144	34.0 (86)	145	16.6 (37)	76	* *
Warrick	447	60.6 (56)	36.2 (62)		609	44.7 (37)	607	14.5 (61)	323	87.6 (44)
Washington	370	43.2 (81)	51.1 (81)		460	58.5 (4)	463	15.1 (53)	221	86.9 (50)
Wayne	1,205	51.1 (74)	42.3 (71)		1,530	53.1 (13)	1,443	13.3 (71)	734	75.1 (79)
Wells	258	61.2 (52)	34.9 (58)		361	44.9 (35)	353	17.3 (31)	179	92.2 (6)
White	395	70.6 (23)	25.3 (27)		550	45.6 (32)	550	13.6 (68)	284	84.2 (63)
Whitley	337	62.0 (50)	34.7 (57)		479	46.6 (28)	475	26.1 (3)	221	90.0 (25)
Unknown Counties	1	* *	* *	* *	305	16.7 (91)	294	17.0 (34)	296	88.5 (40)
Indiana	80,115	63.3	30.0		110,014	40.7	107,563	15.6	55,435	87.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams	619	34.6 (36)	599	21.7 (16)	206	36.4 (16)	497	43.7 (34)	590	39.8 (39)
Allen	8,643	26.9 (8)	8,380	18.2 (8)	2,219	32.9 (34)	7,005	35.1 (15)	8,263	32.4 (23)
Bartholome	1,632	35.8 (42)	1,590	25.3 (37)	564	35.6 (20)	1,289	51.6 (55)	1,570	46.5 (54)
Benton	122	32.8 (23)	118	28.0 (54)	40	* *	96	* *	116	48.3 (59)
Blackford	293	42.3 (75)	289	31.8 (77)	123	29.3 (55)	238	47.9 (41)	286	43.7 (44)
Boone	612	39.1 (55)	586	29.2 (63)	223	33.2 (33)	464	51.3 (53)	579	48.4 (60)
Brown	209	34.4 (32)	202	19.3 (11)	69	* *	166	50.0 (48)	200	44.0 (45)
Carroll	248	33.1 (26)	246	28.0 (54)	82	* *	199	31.7 (12)	243	14.8 (6)
Cass	1,148	25.3 (6)	1,082	18.4 (9)	269	29.7 (52)	917	27.3 (6)	1,067	16.5 (7)
Clark	2,271	34.3 (30)	2,188	23.9 (28)	742	35.0 (23)	1,783	41.4 (26)	2,157	37.1 (32)
Clay	703	39.4 (60)	684	27.2 (49)	267	33.3 (32)	590	56.6 (76)	676	52.7 (77)
Clinton	901	28.0 (11)	882	19.7 (13)	246	34.6 (25)	754	23.2 (3)	870	10.8 (4)
Crawford	219	42.9 (78)	212	38.7 (91)	90	* *	175	64.0 (88)	209	52.6 (76)
Daviess	555	33.9 (29)	514	22.0 (18)	167	34.1 (29)	446	44.4 (35)	502	42.4 (42)
Dearborn	957	45.5 (85)	922	30.8 (72)	412	36.9 (12)	754	53.7 (64)	903	48.7 (62)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Decatur	555	39.8 (62)	540	30.4 (69)	212	26.4 (68)	434	60.6 (82)	536	57.3 (86)
DeKalb	986	38.3 (52)	953	31.6 (76)	362	25.4 (71)	790	58.7 (80)	943	53.4 (79)
Delaware	2,620	34.5 (33)	2,515	28.3 (57)	862	28.4 (60)	1,892	35.3 (16)	2,486	23.3 (12)
Dubois	782	26.6 (7)	767	18.6 (10)	203	34.5 (26)	578	41.0 (24)	757	37.8 (33)
Elkhart	6,105	23.7 (4)	5,895	14.8 (4)	1,376	32.8 (35)	5,122	21.8 (2)	5,832	7.8 (2)
Fayette	649	39.9 (63)	631	29.5 (64)	249	36.1 (19)	524	59.2 (81)	626	55.9 (82)
Floyd	1,409	38.2 (49)	1,342	25.1 (35)	508	40.0 (6)	1,093	48.9 (46)	1,313	46.0 (51)
Fountain	356	46.9 (88)	343	31.5 (75)	162	36.4 (16)	285	61.4 (83)	339	57.5 (87)
Franklin	325	39.1 (55)	316	28.5 (58)	124	37.9 (11)	249	55.0 (69)	310	52.3 (72)
Fulton	517	35.6 (40)	491	24.8 (33)	175	36.6 (14)	422	38.9 (19)	486	33.5 (24)
Gibson	577	41.6 (73)	549	34.2 (86)	222	28.8 (59)	441	57.8 (78)	545	54.3 (80)
Grant	1,756	34.5 (33)	1,684	28.2 (56)	572	25.5 (69)	1,424	42.6 (30)	1,667	23.9 (14)
Greene	521	39.3 (59)	509	28.5 (58)	196	29.1 (57)	387	55.8 (72)	506	45.7 (49)
Hamilton	2,045	22.2 (3)	1,984	14.0 (2)	435	39.8 (7)	1,539	29.9 (10)	1,959	23.3 (12)
Hancock	866	34.5 (33)	834	22.7 (22)	289	34.3 (28)	650	34.5 (13)	823	16.8 (8)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harrison	727	40.0 (64)	694	26.5 (40)	278	39.2 (9)	565	52.2 (58)	682	45.3 (48)
Hendricks	1,273	31.3 (16)	1,136	15.7 (6)	324	46.3 (2)	943	43.3 (32)	1,114	39.4 (37)
Henry	1,003	46.1 (86)	979	31.8 (77)	447	30.6 (46)	827	53.0 (60)	966	49.5 (64)
Howard	2,164	35.2 (39)	2,106	27.0 (46)	738	25.5 (69)	1,762	44.6 (37)	2,076	36.6 (31)
Huntington	726	49.9 (92)	702	31.2 (74)	344	30.8 (44)	621	50.6 (52)	694	46.3 (52)
Jackson	1,070	36.3 (44)	1,046	22.9 (23)	375	38.7 (10)	895	39.1 (22)	1,040	34.6 (28)
Jasper	697	32.6 (22)	652	27.5 (51)	209	21.5 (78)	539	57.5 (77)	648	52.3 (72)
Jay	465	35.1 (38)	450	25.8 (39)	156	29.5 (54)	360	42.5 (29)	440	36.4 (30)
Jefferson	733	42.0 (74)	704	34.7 (89)	293	27.0 (64)	581	40.6 (23)	694	28.4 (19)
Jennings	864	37.4 (48)	840	30.6 (70)	310	15.5 (79)	756	56.0 (73)	829	50.9 (68)
Johnson	2,227	30.8 (14)	2,156	23.1 (24)	649	32.5 (38)	1,624	41.9 (28)	2,116	34.2 (26)
Knox	740	38.2 (49)	654	23.7 (25)	247	30.8 (44)	608	43.4 (33)	634	38.3 (35)
Kosciusko	1,275	30.8 (14)	1,205	22.0 (18)	356	32.3 (39)	1,000	44.4 (35)	1,190	34.1 (25)
LaGrange	434	31.6 (18)	412	23.8 (26)	125	30.4 (47)	337	48.4 (45)	400	42.5 (43)
Lake	13,286	18.1 (2)	12,341	11.6 (1)	2,176	46.2 (3)	9,431	27.8 (8)	12,191	24.6 (16)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
LaPorte	2,734	31.7 (19)	2,567	24.4 (30)	790	31.8 (42)	2,146	43.1 (31)	2,526	39.4 (37)
Lawrence	931	40.5 (69)	901	27.7 (52)	363	36.6 (14)	745	50.3 (50)	895	46.4 (53)
Madison	2,920	34.9 (37)	2,843	27.1 (47)	981	32.0 (41)	2,171	50.3 (50)	2,814	46.8 (55)
Marion	29,025	18.0 (1)	27,471	14.6 (3)	4,769	36.7 (13)	19,086	28.1 (9)	26,505	27.0 (18)
Marshall	1,063	32.9 (24)	1,029	21.9 (17)	331	34.4 (27)	857	20.1 (1)	1,020	5.2 (1)
Martin	172	43.0 (79)	166	30.7 (71)	68	* *	143	41.3 (25)	165	25.5 (17)
Miami	670	40.0 (64)	637	24.5 (31)	247	39.3 (8)	555	54.4 (67)	634	45.9 (50)
Monroe	2,023	34.3 (30)	1,919	21.2 (15)	644	33.7 (31)	1,506	30.6 (11)	1,889	17.5 (10)
Montgomery	872	41.5 (72)	843	31.1 (73)	350	28.9 (58)	718	56.4 (75)	834	49.2 (63)
Morgan	1,275	38.4 (53)	1,228	33.6 (85)	462	26.6 (67)	917	56.3 (74)	1,214	55.0 (81)
Newton	185	27.6 (9)	180	30.0 (67)	47	* *	141	66.7 (90)	180	58.9 (90)
Noble	886	30.0 (12)	846	26.8 (43)	249	22.9 (76)	700	39.0 (20)	839	30.9 (22)
Ohio	29	* *	29	* *	10	* *	22	* *	27	* *
Orange	487	42.3 (75)	471	29.5 (64)	195	26.7 (65)	411	41.6 (27)	467	19.5 (11)
Owen	388	47.2 (89)	377	32.4 (80)	177	28.2 (61)	333	58.0 (79)	372	50.5 (67)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Among women who smoked 3 months prior to pregnancy.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Parke	240	44.2 (80)	240	28.8 (61)	105	35.2 (22)	198	54.0 (66)	237	51.9 (71)
Perry	386	44.6 (82)	374	38.0 (90)	161	24.8 (73)	289	61.9 (85)	368	59.0 (91)
Pike	226	31.9 (20)	216	24.1 (29)	71	* *	174	63.2 (86)	214	57.0 (85)
Porter	2,324	33.2 (27)	2,233	19.6 (12)	736	45.5 (4)	1,736	37.1 (17)	2,190	29.7 (20)
Posey	293	46.4 (87)	275	33.1 (83)	125	29.6 (53)	226	67.3 (91)	273	56.4 (83)
Pulaski	264	33.7 (28)	243	32.5 (81)	81	* *	197	51.3 (53)	240	45.0 (47)
Putnam	732	40.8 (71)	706	26.9 (45)	285	35.4 (21)	601	52.1 (57)	700	48.1 (57)
Randolph	622	33.0 (25)	605	25.6 (38)	196	25.0 (72)	498	45.0 (38)	584	37.8 (33)
Ripley	610	36.9 (47)	582	32.3 (79)	209	24.4 (74)	468	53.2 (61)	569	52.4 (75)
Rush	312	40.1 (67)	305	29.8 (66)	120	30.0 (50)	253	61.7 (84)	302	56.6 (84)
Scott	720	45.3 (84)	697	34.6 (88)	309	27.2 (63)	595	55.6 (71)	690	50.4 (66)
Shelby	929	36.1 (43)	894	26.8 (43)	310	31.3 (43)	706	52.7 (59)	885	51.3 (69)
Spencer	285	36.5 (46)	281	23.8 (26)	103	34.0 (30)	222	46.8 (40)	274	48.2 (58)
St Joseph	5,934	27.9 (10)	5,710	15.1 (5)	1,568	47.8 (1)	4,701	24.7 (4)	5,637	17.2 (9)
Starke	566	40.1 (67)	550	27.1 (47)	217	32.7 (36)	464	53.9 (65)	547	51.4 (70)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Steuben	838	40.0 (64)	816	26.5 (40)	324	30.2 (48)	701	54.8 (68)	802	49.8 (65)
Sullivan	442	42.8 (77)	419	33.4 (84)	172	22.7 (77)	381	53.5 (63)	417	47.5 (56)
Switzerlan	148	49.3 (91)	141	32.6 (82)	68	* *	118	53.4 (62)	138	48.6 (61)
Tippecanoe	3,948	24.4 (5)	3,805	18.0 (7)	917	26.7 (65)	3,180	34.8 (14)	3,754	30.8 (21)
Tipton	182	31.9 (20)	175	30.3 (68)	54	* *	129	49.6 (47)	174	35.6 (29)
Union	143	31.5 (17)	136	25.0 (34)	42	* *	107	55.1 (70)	132	52.3 (72)
Vanderburg	3,077	38.6 (54)	2,690	20.7 (14)	995	41.7 (5)	2,569	26.7 (5)	2,640	9.3 (3)
Vermillion	303	44.6 (82)	297	34.3 (87)	131	28.2 (61)	248	64.5 (89)	294	58.8 (89)
Vigo	2,083	39.4 (60)	1,819	24.6 (32)	681	36.3 (18)	1,747	46.3 (39)	1,802	42.1 (41)
Wabash	634	38.2 (49)	595	27.2 (49)	221	32.6 (37)	510	39.0 (20)	586	23.9 (14)
Warren	185	48.1 (90)	180	26.7 (42)	85	* *	144	63.9 (87)	180	58.3 (88)
Warrick	777	40.7 (70)	744	27.7 (52)	302	30.1 (49)	607	51.7 (56)	737	53.2 (78)
Washington	544	44.3 (81)	530	29.1 (62)	236	32.2 (40)	463	47.9 (41)	521	34.4 (27)
Wayne	1,838	36.4 (45)	1,783	25.2 (36)	643	29.2 (56)	1,524	48.2 (43)	1,760	39.3 (36)
Wells	460	39.1 (55)	449	28.7 (60)	174	29.9 (51)	354	48.3 (44)	444	41.4 (40)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Among women who smoked 3 months prior to pregnancy.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Indiana
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
White	668	30.1 (13)	650	22.2 (20)	191	23.0 (75)	547	27.6 (7)	641	12.5 (5)
Whitley	604	35.6 (40)	585	22.2 (20)	206	35.0 (23)	474	50.2 (49)	583	44.4 (46)
Unknown Counties	307	39.1 (55)	4	* *	1	* *	291	38.5 (18)	3	* *
Indiana	142,169	29.0	135,160	20.5	38,415	34.2	107,925	37.5	132,772	31.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams	572	7.7 (50)	5.2 (18)	467	12.2 (22)	410	2.4 (31)	572	78.8 (3)
Allen	8,042	9.1 (75)	4.9 (13)	6,630	16.4 (81)	5,525	3.5 (72)	8,038	61.4 (59)
Bartholome	1,542	6.8 (22)	6.5 (41)	1,242	13.8 (51)	1,069	2.1 (26)	1,542	68.4 (29)
Benton	112	9.8 (84)	7.1 (60)	87	* *	69	* *	112	64.3 (49)
Blackford	277	7.6 (46)	5.8 (31)	229	11.8 (17)	202	1.5 (13)	277	69.7 (28)
Boone	568	8.1 (58)	6.2 (38)	435	12.2 (22)	382	2.9 (53)	568	56.7 (76)
Brown	192	6.3 (13)	10.9 (91)	158	12.7 (33)	137	0.0 (1)	194	80.9 (1)
Carroll	236	8.5 (65)	9.3 (88)	190	14.7 (65)	162	3.1 (61)	236	72.9 (15)
Cass	1,032	6.6 (18)	6.5 (41)	838	9.8 (2)	748	2.4 (31)	1,038	60.9 (61)
Clark	2,099	8.4 (63)	5.6 (26)	1,672	13.6 (47)	1,437	3.7 (77)	2,097	67.8 (32)
Clay	655	4.9 (2)	7.0 (58)	553	10.1 (3)	497	1.4 (7)	654	62.5 (53)
Clinton	853	6.2 (10)	9.3 (88)	721	12.2 (22)	632	1.9 (21)	852	72.4 (17)
Crawford	202	7.4 (41)	5.4 (21)	164	14.0 (56)	141	1.4 (7)	201	67.7 (34)
Daviess	479	7.1 (30)	2.9 (1)	394	15.7 (74)	328	3.0 (56)	485	58.4 (72)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)	Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Dearborn	880	7.5 (44)	8.9 (83)	718	11.8 (17)	624	2.4 (31)	892	61.1 (60)
Decatur	528	7.6 (46)	5.7 (29)	420	13.8 (51)	362	3.3 (66)	528	66.5 (40)
DeKalb	904	6.6 (18)	8.3 (77)	730	12.6 (32)	635	3.0 (56)	904	77.5 (5)
Delaware	2,435	9.6 (80)	4.6 (10)	1,848	13.7 (49)	1,586	4.1 (80)	2,447	60.2 (63)
Dubois	737	6.5 (17)	6.8 (51)	570	13.2 (39)	493	1.4 (7)	737	75.4 (9)
Elkhart	5,567	7.3 (35)	7.9 (72)	4,751	11.6 (15)	4,182	2.3 (29)	5,586	74.6 (10)
Fayette	613	6.9 (24)	7.0 (58)	495	15.2 (72)	416	1.9 (21)	617	56.2 (78)
Floyd	1,297	9.6 (80)	4.9 (13)	1,037	19.1 (88)	833	3.0 (56)	1,302	65.7 (42)
Fountain	329	7.9 (54)	6.7 (48)	271	14.0 (56)	232	2.2 (28)	331	61.6 (57)
Franklin	304	5.6 (4)	8.2 (76)	237	10.5 (7)	212	2.8 (50)	304	59.2 (69)
Fulton	474	7.4 (41)	5.7 (29)	396	15.9 (78)	328	1.5 (13)	475	66.5 (40)
Gibson	532	6.8 (22)	3.8 (3)	415	13.3 (43)	358	3.1 (61)	537	52.1 (88)
Grant	1,619	9.9 (88)	5.5 (23)	1,321	15.0 (70)	1,119	4.4 (85)	1,621	53.0 (86)
Greene	488	7.2 (32)	6.6 (46)	391	12.5 (29)	341	4.1 (80)	489	59.5 (66)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)		Number	%	Number	%	Number	%	
	Number	% (Rank)	% (Rank)							Number
Hamilton	1,923	8.4 (63)	7.3 (62)	1,460	13.2 (39)	1,263	3.2 (64)	1,930	75.6 (7)	
Hancock	808	6.1 (8)	7.4 (64)	614	12.9 (36)	535	1.1 (6)	808	71.5 (19)	
Harrison	662	8.9 (70)	6.9 (55)	514	16.7 (83)	426	3.5 (72)	662	69.9 (27)	
Hendricks	1,107	7.6 (46)	7.8 (71)	806	11.2 (12)	716	2.7 (47)	1,107	74.3 (11)	
Henry	958	7.8 (52)	7.6 (67)	804	10.1 (3)	721	4.3 (83)	960	68.1 (31)	
Howard	2,017	7.2 (32)	5.9 (32)	1,680	11.6 (15)	1,478	2.4 (31)	2,024	53.7 (84)	
Huntington	673	7.3 (35)	4.8 (12)	581	15.0 (70)	493	3.0 (56)	674	63.9 (50)	
Jackson	1,002	5.9 (5)	6.9 (55)	855	13.8 (51)	731	1.4 (7)	1,004	67.6 (36)	
Jasper	633	7.3 (35)	8.7 (81)	477	12.2 (22)	418	2.6 (42)	634	61.5 (58)	
Jay	430	8.6 (67)	3.7 (2)	343	13.4 (46)	297	4.4 (85)	429	67.6 (36)	
Jefferson	672	9.8 (84)	7.9 (72)	549	16.4 (81)	455	2.6 (42)	673	67.8 (32)	
Jennings	791	7.0 (27)	7.6 (67)	705	14.6 (64)	601	1.8 (19)	792	61.7 (55)	
Johnson	2,073	7.3 (35)	7.6 (67)	1,535	13.3 (43)	1,324	2.6 (42)	2,080	70.2 (25)	
Knox	615	9.1 (75)	4.9 (13)	516	14.5 (62)	441	4.5 (87)	618	55.7 (79)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kosciusko	1,161	7.4 (41)	6.5 (41)	912	14.0 (56)	784	1.4 (7)	1,162	70.3 (23)
LaGrange	392	8.7 (69)	5.9 (32)	313	16.3 (80)	260	2.3 (29)	394	73.4 (13)
Lake	11,914	9.2 (78)	5.6 (26)	8,373	15.8 (75)	7,025	3.0 (56)	11,942	68.4 (29)
LaPorte	2,464	9.1 (75)	5.5 (23)	1,941	14.5 (62)	1,655	3.3 (66)	2,463	60.9 (61)
Lawrence	872	9.7 (83)	5.3 (19)	703	10.4 (6)	627	4.6 (89)	874	65.4 (45)
Madison	2,782	8.1 (58)	5.5 (23)	2,079	13.8 (51)	1,783	3.3 (66)	2,784	57.3 (75)
Marion	26,744	9.0 (72)	6.1 (36)	18,478	14.8 (68)	15,683	2.8 (50)	26,826	63.1 (52)
Marshall	995	6.4 (14)	8.1 (74)	808	11.0 (10)	719	1.9 (21)	995	72.7 (16)
Martin	161	6.2 (10)	7.5 (65)	132	19.7 (89)	106	0.0 (1)	161	54.7 (81)
Miami	605	7.9 (54)	8.1 (74)	514	13.2 (39)	445	1.6 (15)	605	62.0 (54)
Monroe	1,833	7.3 (35)	9.1 (85)	1,488	11.3 (14)	1,319	2.1 (26)	1,834	78.8 (3)
Montgomery	804	7.3 (35)	8.7 (81)	669	12.3 (26)	586	2.6 (42)	805	65.6 (44)
Morgan	1,198	8.0 (56)	6.7 (48)	872	12.5 (29)	762	2.4 (31)	1,200	67.4 (38)
Newton	180	6.1 (8)	6.7 (48)	133	17.3 (85)	110	4.5 (87)	179	57.5 (74)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Noble	819	7.2 (32)	6.5	(41)	653	15.8 (75)	549	3.6 (74)	820	73.4 (13)
Ohio	26	* *	* *	*	22	* *	18	* *	27	* *
Orange	449	8.2 (62)	5.6	(26)	380	13.7 (49)	328	3.4 (69)	449	71.0 (21)
Owen	364	7.7 (50)	8.5	(78)	316	14.9 (69)	268	3.7 (77)	364	70.6 (22)
Parke	235	8.1 (58)	3.8	(3)	197	15.2 (72)	165	3.6 (74)	237	59.1 (70)
Perry	366	10.7 (90)	4.6	(10)	278	14.7 (65)	237	3.4 (69)	366	51.1 (89)
Pike	210	10.0 (89)	7.6	(67)	167	19.8 (90)	131	3.1 (61)	213	53.5 (85)
Porter	2,139	6.4 (14)	8.6	(80)	1,617	11.9 (19)	1,417	2.0 (24)	2,149	71.4 (20)
Posey	266	9.8 (84)	4.5	(8)	208	14.4 (60)	176	3.4 (69)	269	56.5 (77)
Pulaski	235	6.0 (7)	6.8	(51)	173	13.9 (55)	148	2.0 (24)	236	58.9 (71)
Putnam	675	7.1 (30)	8.9	(83)	565	12.7 (33)	487	2.5 (38)	681	64.9 (47)
Randolph	564	9.0 (72)	5.3	(19)	472	13.6 (47)	407	1.7 (17)	567	59.4 (68)
Ripley	555	8.1 (58)	5.9	(32)	434	12.7 (33)	379	1.8 (19)	555	58.4 (72)
Rush	290	6.9 (24)	6.6	(46)	241	12.0 (20)	210	2.9 (53)	292	53.8 (83)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Scott	663	8.9 (70)	4.4 (7)	552	15.8 (75)	463	4.1 (80)	664	52.3 (87)
Shelby	870	7.0 (27)	6.8 (51)	663	10.9 (9)	585	4.3 (83)	876	65.2 (46)
Spencer	268	7.8 (52)	4.5 (8)	211	18.0 (86)	172	1.7 (17)	269	50.2 (90)
St Joseph	5,517	8.5 (65)	7.3 (62)	4,445	12.4 (27)	3,885	2.5 (38)	5,528	73.6 (12)
Starke	528	6.4 (14)	9.1 (85)	435	12.9 (36)	379	3.2 (64)	525	66.7 (39)
Steuben	751	7.6 (46)	9.1 (85)	662	12.1 (21)	573	2.8 (50)	762	80.1 (2)
Sullivan	410	6.6 (18)	6.1 (36)	354	12.4 (27)	310	1.6 (15)	410	70.0 (26)
Switzerlan	132	2.3 (1)	6.8 (51)	108	5.6 (1)	100	0.0 (1)	134	47.8 (91)
Tippecanoe	3,621	6.9 (24)	8.5 (78)	2,997	11.0 (10)	2,656	2.6 (42)	3,630	75.5 (8)
Tipton	166	9.0 (72)	4.2 (5)	117	10.3 (5)	104	3.8 (79)	167	64.7 (48)
Union	130	6.2 (10)	10.8 (90)	100	14.0 (56)	86	* *	129	55.0 (80)
Vanderburg	2,593	9.3 (79)	5.4 (21)	2,207	16.1 (79)	1,834	2.7 (47)	2,600	59.5 (66)
Vermillion	286	8.0 (56)	5.9 (32)	243	14.4 (60)	207	2.4 (31)	286	54.5 (82)
Vigo	1,750	7.0 (27)	7.5 (65)	1,467	13.2 (39)	1,256	2.5 (38)	1,766	63.6 (51)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wabash	573	5.9 (5)	6.3 (39)		469	10.7 (8)	418	1.4 (7)	575	65.7 (42)
Warren	175	5.1 (3)	6.9 (55)		135	13.3 (43)	117	0.9 (4)	175	61.7 (55)
Warrick	712	8.6 (67)	6.5 (41)		558	17.2 (84)	459	0.9 (4)	714	60.1 (64)
Washington	491	9.8 (84)	4.9 (13)		422	14.7 (65)	359	3.6 (74)	492	70.3 (23)
Wayne	1,703	7.5 (44)	6.4 (40)		1,428	12.5 (29)	1,247	2.4 (31)	1,703	60.0 (65)
Wells	429	9.6 (80)	4.2 (5)		337	13.1 (38)	293	2.7 (47)	430	72.1 (18)
White	608	6.6 (18)	7.1 (60)		509	11.2 (12)	447	2.9 (53)	613	67.7 (34)
Whitley	568	10.9 (91)	5.1 (17)		450	18.7 (87)	366	2.5 (38)	567	76.5 (6)
Unknown Counties	3	*	*	*	*	*	0	*	*	*
Indiana	130,177	8.3	6.4		100,826	13.9	86,459	2.8	130,498	65.8

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

