



Forests of Indiana, 2014

Overview

This resource update provides an overview of forest resources in Indiana based on an inventory conducted by the U.S. Forest Service, Forest Inventory and Analysis (FIA) program at the Northern Research Station in cooperation with the Indiana Department of Natural Resources. Estimates are based on field data collected using the FIA annualized strategic sample design and are updated yearly. The estimates presented in this update are for the measurement years 2009-2014 with comparisons between estimates reported in 2009. The current, 2009-2014 sample set consists of 6,492 plots with 1,661 of them being forested or partially forested. About 14.3 percent of the plots were measured each year with observations collected across a period of 6 years. Data used in this publication were accessed from the FIA Database in May 2015. See Bechtold and Patterson (2005) and O’Connell et al. (2014) for definitions and technical details. FIA estimates, tabular data, and maps may be generated at <http://fiatools.fs.fed.us>.

Currently, Indiana houses nearly 4.9 million acres of forest land. Forested area has increased by about 3.5 percent (166,000 acres) since 2009 (Table 1). Timberland accounts for nearly 97 percent, while the remaining 3 percent of forest is reserved or unproductive. The number of live trees on Indiana’s forest land in 2014 was estimated at 2.2 billion trees, an increase of 2.5 percent from 2009. Statewide, the number of trees, one inch and larger per acre in 2014 averages 452. Net volume (10.5 billion ft³) experienced an increase of about 8.4 percent, which statewide in 2014, averages 2.1 thousand cubic feet per acre (statewide, about 26 cords per acre). Live tree biomass (above ground) is estimated at 272.3 million oven-dry tons or statewide, on average, about 55.8 tons per acre. Average annual net growth decreased by 27.5 percent partly due to reversions of nonforest to forest¹. Statewide, average annual net growth is 46 cubic feet per acre per year while annual harvest removals decreased by 18.8 percent. Annual mortality increased by 30.4 percent between 2009 and 2014; however, when put in perspective as a percentage of net volume, mortality in 2009 was 1.0 percent of net volume and mortality in 2014 was 1.24 percent of net volume, a difference of only 0.24 percent from 2009. Similar trends were observed on Indiana’s timberlands (Table 1).

Table 1.—Indiana forest statistics, change between 2009 and 2014

	2009 Estimate	Sampling error (percent)	2014 Estimate	Sampling error (percent)	Percent change since 2009
Forest Land					
Area (thousand acres)	4,713.6	1.1	4,879.6	1.0	3.5
Number of live trees ≥ 1 in diameter (million trees)	2,154	2.0	2,208.1	2.0	2.5
Net volume live trees ≥ 5 in diameter (million ft ³)	9,695.8	1.7	10,511.2	1.7	8.4
Live tree aboveground biomass (thousand oven-dry tons)	254,240.3	1.6	272,327.0	1.5	7.1
Net growth live trees ≥ 5 in (thousand ft ³ /yr)	316,065.6	5.1	229,165.2	4.7	-27.5
Harvest removals of live trees ≥ 5 in (thousand ft ³ /yr)	90,920.1	14.5	73,784.9	14.2	-18.8
Annual mortality of live trees ≥ 5 in (thousand ft ³ /yr)	100,300.4	7.6	130,776.8	6.3	30.4
Timberland					
Area (thousand acres)	4,568.1	1.2	4,717.1	1.1	3.3
Number of live trees ≥ 1 in diameter (million trees)	2,071.6	2.1	2,116.2	2.1	2.2
Net volume live trees ≥ 5 in diameter (million ft ³ /yr)	9,380.9	1.8	10,149.8	1.8	8.2
Live tree aboveground biomass (thousand oven-dry tons)	245,858.8	1.9	262,733.6	1.6	6.9
Net growth of growing-stock trees ≥ 5 in (thousand ft ³ /yr)	272,017.1	5.1	213,073.2	4.4	-21.7
Harvest removals of growing-stock trees ≥ 5 in (thousand ft ³ /yr)	82,617.0	15.2	65,961.7	14.5	-20.2
Annual mortality of growing-stock trees ≥ 5 in (thousand ft ³ /yr)	76,470.4	8.9	92,649.0	7.4	21.2

¹ Some nonforest to forest reversions that occurred during and prior to the 1990’s to early 2000’s were not identified until early on in the adoption of the annual inventory system and stem from use of enhanced imagery and GIS technology that began in the early 2000’s. We improved our ability to detect forest early on in the implementation of the annual inventory system and thus, more plots were field measured. As a result, higher estimates of growth were recorded in earlier annual inventories resulting in an artificially inflated growth estimate.



Forest Area

Indiana is divided into four survey units, with forest land (4.9 million acres) unevenly distributed among units: Northern (1.4 million acres), Lower Wabash (977 thousand acres), Upland Flats (677 thousand acres), and Knobs (1.83 million acres (Fig. 1). The three southern tier survey units comprise about 40 percent of the land and water area but contain over 70 percent of the forest with the Knobs survey unit containing about 40 percent of the forest.

Eighty four percent or over 4.1 million acres of forest land is privately owned. The state and local government owns 8.0 percent or 391 thousand acres of forest land while the federal government owns roughly 7.4 percent or 380 thousand acres. A little over 3 percent or 161 thousand acres of forest land is considered reserved.




Figure 1.—Forest land (dark green) by survey unit, Indiana.

The total area of Indiana’s forest land (4.9 million acres) and timberland (4.7 million acres) has modest increases over the past several decades following a trend since 1967 (Fig. 2).




Figure 2.—Area of forest land and timberland in Indiana by inventory year, 1950, 1965, 1986, 1998, 2003 to 2014. Error bars represent 1 standard error or a 67-percent confidence interval.

Hardwoods are the dominant species in Indiana. Some forest-type groups are much more common than others. The oak/hickory group alone occupies 71 percent of forest land, the bulk of which resides in the white oak/red oak/hickory forest type (1.5 million acres). Softwoods alone occupy 108 thousand acres. The oak/pine group occupies almost 152,500 acres which represents 3 percent of the forest land wood basket.

Forest land consists mainly of sawtimber stands (78 percent); 14 percent of forest land is made up of poletimber stands, 7 percent contain seedling-sapling stands and less than 1 percent is considered nonstocked.

Indicative of a maturing (aging) forest, white and red oak/hickory is found primarily in the large stand-size class (Fig. 3). The cherry/white ash/yellow -poplar group is less common (468 thousand acres) as are the mixed upland hardwoods (406 thousand acres). Both show similar distributions across stand-size classes with a large proportion in the medium and large diameter classes. The sugar maple/beech/yellow birch forest-type group is relatively abundant (224 thousand acres) and occurs mostly in large stand-size classes (Fig. 3).

Currently, nearly half (48 percent) of the stands are over 61 years of age.




Figure 3.—Area of forest land by five most common forest-type groups and stand-size classes, Indiana, 2014. Error bars represent 1 standard error or a 67-percent confidence interval.

Volume, Biomass, and Trends

Crews recorded 95 species (including unknowns collected to the genus level) on Indiana forest land in the measurement years included in the 2014 dataset. Hardwoods are the dominate forest land species comprising 97 percent. Sugar maple (*Acer saccharum*) is by far the most numerous tree in Indiana with an estimated 360 million individuals; red maple (*Acer rubrum*) is second behind with an estimated 110 million trees in Indiana (Table 2).

Interestingly, the most numerous species, sugar maple, is not the most voluminous species in the state. That distinction belongs to the State tree, yellow-poplar, a.k.a. the tulip tree (*Liriodendron tulipifera*) with a net volume of nearly 1.24 billion cubic feet. These yellow-poplars also store approximately 23.5 million tons of woody biomass in their tissues.

Of the top 10 most voluminous species, yellow-poplar and sugar maple are growing the most vigorously with each accumulating over 37 and 26 million cubic feet per year respectively.

Black oak followed by yellow-poplar, white ash, and sugar maple show the highest mortality in Indiana. Black oaks that are stressed from drought, gypsy moth defoliation, old age, fire, poor site conditions, or other factors often succumb to secondary agents such as twolined chestnut borer (*Agilus bilineatus*), Hypoxylon canker (*Hypoxylon mammatum*), and shoestring root rot (*Armillaria mellea*). This scenario, in which a primary agent stresses the tree and a secondary agent kills it, is known as "oak decline" and is responsible for considerable black oak mortality.

Several species are removed in harvests, but yellow-poplar removals are nearly double or more, by volume, than most species except sugar maple, white ash, and black oak (Table 2).

Biomass is distributed throughout the State, with the largest concentrations in the southern tier of Indiana (Fig. 4)




Figure 4.—Distribution of live-tree and sapling biomass on forest land, Indiana 2014

Table 2.—Number, volume, biomass, growth, mortality, and removals of live trees on forest land by species of the top 10 tree species by net volume, Indiana, 2009-2014.

Common Name	Latin Name	Million Trees ^a	Net Volume ^b (million ft ³)	Aboveground Biomass ^a (thousand dry tons)	Average Annual Net Growth ^b (thousand ft ³)	Average Annual Mortality ^b (thousand ft ³)	Average Annual Harvest Removals ^b (thousand ft ³)
Yellow-poplar	<i>Liriodendron tulipifera</i>	74.55	1,238.95	23,525.10	37,160.77	10,249.96	10,999.28
Sugar maple	<i>Acer saccharum</i>	359.91	1,113.08	33,016.16	26,202.99	9,227.39	7,779.98
White oak	<i>Quercus alba</i>	35.98	747.48	20,649.19	12,832.32	3,897.60	4,967.37
White ash	<i>Fraxinus americana</i>	99.26	567.16	15,497.47	10,320.22	10,094.14	6,604.36
Black oak	<i>Quercus velutina</i>	34.12	566.92	15,576.87	6,200.11	11,552.40	7,380.07
Red maple	<i>Acer rubrum</i>	110.16	465.74	11,411.58	13,233.55	3,008.57	1,035.61
Northern red oak	<i>Quercus rubra</i>	24.05	450.12	12,624.32	10,471.73	4,964.90	4,165.83
American sycamore	<i>Platanus occidentalis</i>	16.51	468.63	9,567.00	13,404.59	1,371.43	2,697.38
Shagbark hickory	<i>Carya ovata</i>	44.86	365.26	11,591.81	6,582.19	1,075.14	1,619.30
Black cherry	<i>Prunus serotina</i>	40.75	350.62	11,093.26	6,245.69	1,435.66	1,451.20

^a Trees ≥ 1 in diameter

^b Trees ≥ 5 in diameter

Indiana's Timber Product Output, 2013

Primary wood-using mills in Indiana were surveyed to determine the size and composition of the State's primary wood-using industry, its use of roundwood, and the generation and disposition of wood residues. Below are some preliminary findings from that survey. A full report is in preparation as more detailed information is analyzed.

There were 67.7 million cubic feet of industrial roundwood harvested from Indiana's forests in 2013, an 8 percent increase from 2008. Saw logs accounted for 92 percent of industrial roundwood harvested in 2013 (Fig. 5). Veneer logs, which were the second most harvested product, only accounted for 3 percent of the total harvest.




Figure 5.—Industrial roundwood production by product, Indiana, 2008 and 2013.

Yellow-poplar replaced red oaks as the most harvested species in 2013, making up 20 percent of the total harvest (Fig. 6). Other important species harvested were red and white oaks, ash, and hard maple. Between 2008 and 2013, the harvesting of ash increased by 50 percent.




Figure 6.—Industrial roundwood production by species group, Indiana, 2008 and 2013.

There was 1.1 million green tons of mill residues generated during the processing of industrial roundwood in 2013. Two-thirds of the mill residues were either used by the pulp and paper industry, or used for mulch (Fig. 7). Only one percent of the mill residues from Indiana's primary wood using industry went unused.




Figure 7.—Distribution of residues generated by primary wood-using mills by method of disposal, Indiana, 2013.

References

Bechtold, W.A.; Patterson, P.L., eds. 2005. **The enhanced Forest Inventory and Analysis program: national sampling design and estimation procedures**. Gen. Tech. Rep. SRS-80. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 85 p.

O'Connell, B.M.; LaPoint, E.B.; Turner, J.A.; Ridley, T.; Pugh, S.A.; Wilson, A.M.; Waddell, K.L.; Conkling, B.L. 2014. **The Forest Inventory and Analysis database: database description and users manual version 6.0.1 for Phase 2**. Washington, DC: U.S. Department of Agriculture, Forest Service. (<http://www.fia.fs.fed.us/library/database-documentation/>)

Walters, Brian F.; Settle, Jeff; Piva, Ronald J. 2012. **Indiana timber industry: an assessment of timber product output and use, 2008**. Resour. Bull. NRS-63. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 72 p.

How to Cite This Publication

Gormanson, Dale D.; Piva, Ronald J. 2015. Forest of Indiana, 2014. Resource Update FS-xx. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 4 p.

Contact Information

Dale D. Gormanson, Forester
 USDA Forest Service, Northern Research Station
 1992 Folwell Ave.
 St. Paul, MN 55108
 Ph: 651-649-5126/ Fax:651-649-5140
dgormanson@fs.fed.us
 Northern FIA: <http://nrs.fs.fed.us/fia/>
 National FIA: <http://fia.fs.fed.us>

USDA is an equal opportunity provider and employer

The published report is available online at <http://treesearch.fs.fed.us>