

# Summary Chart of U.S. Medical Eligibility Criteria for Contraceptive Use

Condition	Sub-Condition	Cu-IUD		LNG-IUD		Implant		DMPA		POP		CHC	
		I	C	I	C	I	C	I	C	I	C	I	C
Age		Menarche to <20 yrs: <b>2</b>		Menarche to <20 yrs: <b>2</b>		Menarche to <18 yrs: <b>1</b>		Menarche to <18 yrs: <b>2</b>		Menarche to <18 yrs: <b>1</b>		Menarche to <40 yrs: <b>1</b>	
		<b>≥20 yrs:<b>1</b></b>		<b>≥20 yrs:<b>1</b></b>		<b>18-45 yrs:<b>1</b></b>		<b>18-45 yrs:<b>1</b></b>		<b>18-45 yrs:<b>1</b></b>		<b>≥40 yrs:<b>2</b></b>	
Anatomical abnormalities	a) Distorted uterine cavity	<b>4</b>	<b>4</b>										
	b) Other abnormalities	<b>2</b>	<b>2</b>										
Anemias	a) Thalassemia	<b>2</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
	b) Sickle cell disease <sup>‡</sup>	<b>2</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>2</b>	
Benign ovarian tumors <i>(including cysts)</i>	c) Iron-deficiency anemia	<b>2</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
		<b>1</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
Breast disease	a) Undiagnosed mass	<b>1</b>	<b>2</b>	<b>2*</b>		<b>2*</b>		<b>2*</b>		<b>2*</b>		<b>2*</b>	
	b) Benign breast disease	<b>1</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
Breastfeeding	c) Family history of cancer	<b>1</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
	d) Breast cancer <sup>‡</sup>												
	i) Current	<b>1</b>	<b>4</b>	<b>4</b>		<b>4</b>		<b>4</b>		<b>4</b>		<b>4</b>	
	ii) Past and no evidence of current disease for 5 years	<b>1</b>	<b>3</b>	<b>3</b>		<b>3</b>		<b>3</b>		<b>3</b>		<b>3</b>	
Cervical cancer	a) <21 days postpartum					<b>2*</b>		<b>2*</b>		<b>2*</b>		<b>4*</b>	
	b) 21 to <30 days postpartum												
	i) With other risk factors for VTE					<b>2*</b>		<b>2*</b>		<b>2*</b>		<b>3*</b>	
	ii) Without other risk factors for VTE					<b>2*</b>		<b>2*</b>		<b>2*</b>		<b>3*</b>	
Cervical ectropion	c) 30-42 days postpartum												
	i) With other risk factors for VTE					<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>3*</b>	
Cervical intraepithelial neoplasia	ii) Without other risk factors for VTE					<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>2*</b>	
	d) >42 days postpartum					<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>2*</b>	
Cervical cancer	Awaiting treatment	<b>4</b>	<b>2</b>	<b>4</b>	<b>2</b>	<b>2</b>		<b>2</b>		<b>1</b>		<b>2</b>	
Cervical ectropion		<b>1</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
Cervical intraepithelial neoplasia		<b>1</b>	<b>2</b>	<b>2</b>		<b>2</b>		<b>2</b>		<b>1</b>		<b>2</b>	
Cirrhosis	a) Mild ( <i>compensated</i> )	<b>1</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
	b) Severe <sup>‡</sup> ( <i>decompensated</i> )	<b>1</b>	<b>3</b>	<b>3</b>		<b>3</b>		<b>3</b>		<b>4</b>			
Cystic fibrosis <sup>‡</sup>		<b>1*</b>	<b>1*</b>	<b>1*</b>		<b>2*</b>		<b>1*</b>		<b>1*</b>			
Deep venous thrombosis (DVT)/Pulmonary embolism (PE)	a) History of DVT/PE, not receiving anticoagulant therapy												
	i) Higher risk for recurrent DVT/PE	<b>1</b>	<b>2</b>	<b>2</b>		<b>2</b>		<b>2</b>		<b>4</b>			
	ii) Lower risk for recurrent DVT/PE	<b>1</b>	<b>2</b>	<b>2</b>		<b>2</b>		<b>2</b>		<b>3</b>			
	b) Acute DVT/PE	<b>2</b>	<b>2</b>	<b>2</b>		<b>2</b>		<b>2</b>		<b>4</b>			
	c) DVT/PE and established anticoagulant therapy for at least 3 months												
	i) Higher risk for recurrent DVT/PE	<b>2</b>	<b>2</b>	<b>2</b>		<b>2</b>		<b>2</b>		<b>4*</b>			
	ii) Lower risk for recurrent DVT/PE	<b>2</b>	<b>2</b>	<b>2</b>		<b>2</b>		<b>2</b>		<b>3*</b>			
	d) Family history ( <i>first-degree relatives</i> )	<b>1</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>2</b>			
	e) Major surgery												
	i) With prolonged immobilization	<b>1</b>	<b>2</b>	<b>2</b>		<b>2</b>		<b>2</b>		<b>4</b>			
	ii) Without prolonged immobilization	<b>1</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>2</b>			
	f) Minor surgery without immobilization	<b>1</b>	<b>1</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>			
Depressive disorders		<b>1*</b>	<b>1*</b>	<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>	

**Key:**

**1** No restriction (method can be used)

**3** Theoretical or proven risks usually outweigh the advantages

**2** Advantages generally outweigh theoretical or proven risks

**4** Unacceptable health risk (method not to be used)

Condition	Sub-Condition	Cu-IUD		LNG-IUD		Implant		DMPA		POP		CHC	
		I	C	I	C	I	C	I	C	I	C	I	C
Diabetes	a) History of gestational disease	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
	b) Nonvascular disease												
	i) Non-insulin dependent	<b>1</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>	
	ii) Insulin dependent	<b>1</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>	
	c) Nephropathy/retinopathy/neuropathy <sup>‡</sup>	<b>1</b>		<b>2</b>		<b>2</b>		<b>3</b>		<b>2</b>		<b>3/4*</b>	
	d) Other vascular disease or diabetes of >20 years' duration <sup>‡</sup>	<b>1</b>		<b>2</b>		<b>2</b>		<b>3</b>		<b>2</b>		<b>3/4*</b>	
Dysmenorrhea	Severe	<b>2</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
Endometrial cancer <sup>‡</sup>		<b>4</b>	<b>2</b>	<b>4</b>	<b>2</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
Endometrial hyperplasia		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
Endometriosis		<b>2</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
Epilepsy <sup>‡</sup>	(see also Drug Interactions)	<b>1</b>		<b>1</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>	
Gallbladder disease	a) Symptomatic												
	i) Treated by cholecystectomy	<b>1</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>	
	ii) Medically treated	<b>1</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>3</b>	
	iii) Current	<b>1</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>3</b>	
	b) Asymptomatic	<b>1</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>	
Gestational trophoblastic disease <sup>‡</sup>	a) Suspected GTD (immediate postevacuation)												
	i) Uterine size first trimester	<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>	
	ii) Uterine size second trimester	<b>2*</b>		<b>2*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>	
	b) Confirmed GTD												
	i) Undetectable/non-pregnant β-hCG levels	<b>1*</b>	<b>1*</b>	<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>	
	ii) Decreasing β-hCG levels	<b>2*</b>	<b>1*</b>	<b>2*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>	
	iii) Persistently elevated β-hCG levels or malignant disease, with no evidence or suspicion of intrauterine disease	<b>2*</b>	<b>1*</b>	<b>2*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>	
	iv) Persistently elevated β-hCG levels or malignant disease, with evidence or suspicion of intrauterine disease	<b>4*</b>	<b>2*</b>	<b>4*</b>		<b>2*</b>		<b>1*</b>		<b>1*</b>		<b>1*</b>	
Headaches	a) Nonmigraine (mild or severe)	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
	b) Migraine												
	i) Without aura (includes menstrual migraine)	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>2*</b>	
	ii) With aura	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>4*</b>	
History of bariatric surgery <sup>‡</sup>	a) Restrictive procedures	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
	b) Malabsorptive procedures	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>3</b>	
													<b>COCs: 3</b>
History of cholestasis	a) Pregnancy related	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>2</b>	
	b) Past COC related	<b>1</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>2</b>		<b>3</b>	
History of high blood pressure during pregnancy		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>2</b>	
History of Pelvic surgery		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
HIV	a) High risk for HIV	<b>1*</b>	<b>1*</b>	<b>1*</b>	<b>1*</b>	<b>1</b>		<b>1</b>		<b>1</b>		<b>1</b>	
	b) HIV infection							<b>1*</b>		<b>1*</b>		<b>1*</b>	

# Summary Chart of U.S. Medical Eligibility Criteria for Contraceptive Use

Condition	Sub-Condition	Cu-IUD		LNG-IUD		Implant		DMPA		POP		CHC	
		I	C	I	C	I	C	I	C	I	C	I	C
Hypertension	a) Adequately controlled hypertension	1*		1*		1*		2*		1*		3*	
	b) Elevated blood pressure levels ( <i>properly taken measurements</i> )												
	i) Systolic 140–159 or diastolic 90–99	1*		1*		1*		2*		1*		3*	
	ii) Systolic ≥160 or diastolic ≥100 <sup>‡</sup>	1*		2*		2*		3*		2*		4*	
	c) Vascular disease	1*		2*		2*		3*		2*		4*	
Inflammatory bowel disease	( <i>Ulcerative colitis, Crohn's disease</i> )	1		1		1		2		2		2/3*	
Ischemic heart disease <sup>‡</sup>	Current and history of	1		2	3	2	3	3		2	3	4	
Known thrombogenic mutations <sup>‡</sup>		1*		2*		2*		2*		2*		4*	
Liver tumors	a) Benign												
	i) Focal nodular hyperplasia	1		2		2		2		2		2	
	ii) Hepatocellular adenoma <sup>‡</sup>	1		3		3		3		3		4	
b) Malignant <sup>‡</sup> (hepatoma)		1		3		3		3		3		4	
Malaria		1		1		1		1		1		1	
Multiple risk factors for atherosclerotic cardiovascular disease	(e.g., older age, smoking, diabetes, hypertension, low HDL, high LDL, or high triglyceride levels)	1		2		2*		3*		2*		3/4*	
Multiple sclerosis	a) With prolonged immobility	1		1		1		2		1		3	
	b) Without prolonged immobility	1		1		1		2		1		1	
Obesity	a) Body mass index (BMI) ≥30 kg/m <sup>2</sup>	1		1		1		1		1		2	
	b) Menarche to <18 years and BMI ≥ 30 kg/m <sup>2</sup>	1		1		1		2		1		2	
Ovarian cancer <sup>‡</sup>		1		1		1		1		1		1	
Parity	a) Nulliparous	2		2		1		1		1		1	
	b) Parous	1		1		1		1		1		1	
Past ectopic pregnancy		1		1		1		1		2		1	
Pelvic inflammatory disease	a) Past												
	i) With subsequent pregnancy	1		1		1		1		1		1	
	ii) Without subsequent pregnancy	2		2		2		1		1		1	
Peripartum cardiomyopathy <sup>‡</sup>	b) Current	4		2*	4	2*		1		1		1	
	a) Normal or mildly impaired cardiac function												
	i) <6 months	2		2		1		1		1		4	
	ii) ≥6 months	2		2		1		1		1		3	
	b) Moderately or severely impaired cardiac function	2		2		2		2		2		4	
Postabortion	a) First trimester	1*		1*		1*		1*		1*		1*	
	b) Second trimester	2*		2*		1*		1*		1*		1*	
	c) Immediate postseptic abortion	4		4		1*		1*		1*		1*	
Postpartum (nonbreastfeeding women)	a) <21 days					1		1		1		4	
	b) 21 days to 42 days												
	i) With other risk factors for VTE					1		1		1		3*	
	ii) Without other risk factors for VTE							1		1		2	
	c) >42 days							1		1		1	
Postpartum (in breastfeeding or non-breastfeeding women, including cesarean delivery)	a) <10 minutes after delivery of the placenta												
	i) Breastfeeding	1*		2*									
	ii) Nonbreastfeeding	1*		1*									
	b) 10 minutes after delivery of the placenta to <4 weeks	2*		2*									
	c) ≥4 weeks	1*		1*									
	d) Postpartum sepsis	4		4									

Condition	Sub-Condition	Cu-IUD		LNG-IUD		Implant		DMPA		POP		CHC	
		I	C	I	C	I	C	I	C	I	C	I	C
Pregnancy													
Rheumatoid arthritis	a) On immunosuppressive therapy	2		1		2		1		NA*		NA*	
	b) Not on immunosuppressive therapy	1		1		1		1		2/3*		1	
Schistosomiasis	a) Uncomplicated	1		1		1		1		1		1	
	b) Fibrosis of the liver <sup>‡</sup>	1		1		1		1		1		1	
Sexually transmitted diseases (STDs)	a) Current purulent cervicitis or chlamydial infection or gonococcal infection	4		2*	4	2*		1		1		1	
	b) Vaginitis ( <i>including trichomonas vaginalis and bacterial vaginosis</i> )	2		2		2		1		1		1	
	c) Other factors relating to STDs	2*		2	2*	2		1		1		1	
Smoking	a) Age <35	1		1		1		1		1		1	
	b) Age ≥35, <15 cigarettes/day	1		1		1		1		1		3	
	c) Age ≥35, ≥15 cigarettes/day	1		1		1		1		1		4	
Solid organ transplantation <sup>‡</sup>	a) Complicated	3		2		3		2		2		2	
	b) Uncomplicated	2		2		2		2		2		2	
Stroke <sup>‡</sup>	History of cerebrovascular accident	1		2		2	3	3		2	3	4	
Superficial venous disorders	a) Varicose veins	1		1		1		1		1		1	
	b) Superficial venous thrombosis (acute or history)	1		1		1		1		1		3*	
Systemic lupus erythematosus <sup>‡</sup>	a) Positive (or unknown) antiphospholipid antibodies	1*		1*		3*		3*		3*		3*	
	b) Severe thrombocytopenia	3*		2*		2*		2*		3*		2*	
	c) Immunosuppressive therapy	2*		1*		2*		2*		2*		2*	
	d) None of the above	1*		1*		2*		2*		2*		2*	
Thyroid disorders	Simple goiter/ hyperthyroid/hypothyroid	1		1		1		1		1		1	
Tuberculosis <sup>‡</sup> ( <i>see also Drug Interactions</i> )	a) Nonpelvic	1		1		1		1		1*		1*	
	b) Pelvic	4		3		4	3	1*		1*		1*	
Unexplained vaginal bleeding	(suspicious for serious condition) before evaluation	4*		2*	4*	2*		3*		3*		2*	
Uterine fibroids						2		2		1		1	
Valvular heart disease	a) Uncomplicated					1		1		1		1	
	b) Complicated <sup>‡</sup>					1		1		1		4	
Vaginal bleeding patterns	a) Irregular pattern without heavy bleeding					1		1		2		2	
	b) Heavy or prolonged bleeding					2*		1*	2*	2*		2*	
Viral hepatitis	a) Acute or flare					1		1		1		1	
	b) Carrier/Chronic					1		1		1		3/4*	2
<b>Drug Interactions</b>													
Antiretrovirals used for prevention (PrEP) or treatment of HIV	Fosamprenavir (FPV)			1/2*	1*	1/2*	1*		2*		2*		3*
	All other ARVs are 1 or 2 for all methods.												
Anticonvulsant therapy	a) Certain anticonvulsants (phenytoin, carbamazepine, barbiturates, primidone, topiramate, oxcarbazepine)			1		1			2*		1*		3*
	b) Lamotrigine			1		1			1		1		3*
Antimicrobial therapy	a) Broad spectrum antibiotics			1		1			1		1		1
	b) Antifungals			1		1			1		1		1
SSRIs	c) Antiparasitics			1		1			1		1		1
	d) Rifampin or rifabutin therapy			1		1			2*		1*		3*
St. John's wort				1		1			1		1		2

**Updated in 2020.** This summary sheet only contains a subset of the recommendations from the U.S. MEC. For complete guidance, see: [https://www.cdc.gov/reproductivehealth/contraception/contraception\\_guidance.htm](https://www.cdc.gov/reproductivehealth/contraception/contraception_guidance.htm). Most contraceptive methods do not protect against sexually transmitted diseases (STDs). Consistent and correct use of the male latex condom reduces the risk of STDs and HIV.