

Sensitive Population - FISH CONSUMPTION ADVISORY

Location	Species	Fish Size	Maximum Amount for Adults to Eat
East Fork of Wildcat Creek Howard County	Common Carp	up to 23 23+	1 meal / month (8 ounces / month) Do not eat
Kokomo Creek Howard County	All Fish	ALL	Do not eat
Kokomo Reservoir Howard County	Black Crappie	up to 11	1 meal / week (8 ounces / week)
	Bluegill	up to 7	1 meal / week (8 ounces / week)
	Common Carp	up to 19 19+	1 meal / week (8 ounces / week) 1 meal / month (8 ounces / month)
	Walleye	up to 16	1 meal / week (8 ounces / week)
	White Bass	up to 10	1 meal / week (8 ounces / week)
	White Crappie	up to 10	1 meal / week (8 ounces / week)
Wildcat Creek Howard County Upstream of the Waterworks Dam in Kokomo	Bluegill	up to 6	1 meal / week (8 ounces / week)
	Common Carp	up to 19 19+	1 meal / month (8 ounces / month) Do not eat
	Longear Sunfish	up to 5	1 meal / week (8 ounces / week)
	Rock Bass	up to 7	1 meal / week (8 ounces / week)
Carroll/Howard County Downstream of the Waterworks Dam in Kokomo through Howard and Carroll counties.	All Fish	ALL	Do not eat

Sensitive Population

*Sensitive populations include: pregnant or nursing women, women that will become pregnant, and children under 6 years of age. These consumers should use caution when eating some types of sportfish.

**The maximum amount for adults to eat is determined by fishing location, species and fish length. For instance, if you eat the maximum amount of one species during a week, you should not eat any other fish which has a consumption limitation until the following week. Example - if you eat the limit (4 ounces) of a 20 inch long Carp from the Easy Catch River, then you should not eat any other sportfish from the list until the following week.