

State Prevalence Rankings: BRFSS, 2009

State:	Current Smokers		Ever Told Have High Blood Pressure		Obese Based on BMI		No Moderate Phys. Activity on 5 or More Days in Past Week		Diabetes		High Cholesterol		Current Asthma	
	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank
Alabama	22.5	7	37.2	3	31.6	7	58.9	4	12.2	2	39.9	7	7.6	41
Alaska	20.6	12	26.4	43	25.4	36	39.3	51	5.8	50	34.9	45	9.0	21
Arizona	16.1	40	26.5	42	25.8	32	49.4	24	8.4	24	40.9	4	10.8	2
Arkansas	21.4	11	34.4	6	31.4	8	52.6	12	10.1	9	38.7	16	7.5	44
California	12.8	50	25.6	48	25.5	33	48.6	28	9.1	18	36.5	36	7.8	36
Colorado	17.0	35	22.3	50	18.9	51	42.9	45	5.7	51	35.2	43	8.2	32
Connecticut	15.4	43	27.1	36	21.0	49	46.0	41	6.6	46	37.2	29	9.3	17
Delaware	18.2	23	30.7	16	27.5	24	49.0	26	8.1	30	38.2	23	8.8	25
District of Co.	15.2	45	26.1	44	20.1	50	45.5	42	7.5	38	34.5	47	9.8	12
Florida	18.1	24	30.9	15	25.9	31	52.1	14	9.8	11	39.5	11	7.1	46
Georgia	17.6	30	31.2	14	27.7	23	54.2	9	9.5	13	37.0	31	6.9	47
Hawaii	15.3	44	30.2	18	22.9	47	46.7	37	8.4	24	38.8	13	9.4	16
Idaho	16.3	39	25.8	46	25.1	39	42.4	47	7.9	32	37.2	29	8.3	31
Illinois	18.5	22	28.8	25	27.3	25	48.2	31	8.1	30	37.5	25	9.0	21
Indiana	23.1	5	31.3	13	29.9	14	52.0	15	9.3	14	39.8	9	9.1	18
Iowa	17.1	33	28.0	30	28.4	19	50.3	21	7.6	37	37.5	25	6.8	48
Kansas	17.8	28	28.6	26	28.7	18	51.5	18	8.5	23	38.5	20	8.5	30
Kentucky	25.6	1	36.3	4	32.3	4	54.3	8	11.5	4	41.6	2	10.1	6
Louisiana	22.0	8	35.6	5	33.9	2	56.4	5	11.0	5	36.9	33	6.3	51
Maine	17.2	32	30.0	20	26.4	28	43.7	43	8.3	26	38.8	13	10.8	2
Maryland	15.1	46	29.4	23	26.7	27	51.3	20	9.3	14	37.3	28	9.1	18
Massachusetts	14.9	48	25.7	47	21.8	48	47.0	36	7.9	32	35.6	42	10.8	2
Michigan	19.6	18	29.7	22	30.3	10	48.0	32	9.3	14	38.6	18	10.0	7
Minnesota	16.7	37	21.5	51	25.3	37	47.2	34	6.3	47	33.8	49	6.6	49
Mississippi	23.3	4	37.4	2	35.3	1	62.5	3	11.6	3	41.3	3	7.5	44
Missouri	23.1	5	30.6	17	30.5	9	50.0	22	7.9	32	38.1	24	9.5	15
Montana	16.8	36	27.7	32	23.7	44	41.3	50	6.7	45	36.5	36	8.1	33
Nebraska	16.7	37	27.1	36	28.0	21	48.9	27	7.4	40	37.4	27	7.6	41
Nevada	22.0	8	27.5	34	26.4	28	48.5	30	7.9	32	38.5	20	9.0	21
New Hampshire	15.7	42	28.6	26	26.3	30	46.6	39	7.1	42	38.6	18	10.2	5
New Jersey	15.8	41	28.1	29	23.9	43	52.4	13	8.7	21	37.0	31	7.6	41
New Mexico	17.9	25	26.6	41	25.5	33	46.7	37	8.6	22	34.1	48	8.6	29
New York	17.9	25	28.5	28	24.5	41	49.2	25	8.8	20	38.9	12	9.8	12
North Carolina	20.3	13	31.5	11	30.1	12	53.5	10	9.6	12	39.9	7	7.8	36
North Dakota	18.6	21	26.7	40	28.3	20	47.7	33	7.5	38	34.7	46	8.8	25
Ohio	20.3	13	31.7	10	29.7	15	51.4	19	10.1	9	39.6	10	9.9	9
Oklahoma	25.4	3	34.3	7	32.0	5	52.8	11	11.0	5	40.2	6	9.9	9
Oregon	17.8	28	27.1	36	23.6	45	43.4	44	8.2	27	35.2	43	11.1	1
Pennsylvania	20.2	16	31.4	12	28.0	21	49.5	23	9.0	19	38.8	13	9.1	18
Rhode Island	15.0	47	30.1	19	24.9	40	51.6	17	7.0	43	36.4	38	10.0	7
South Carolina	20.3	13	32.6	8	30.0	13	54.6	7	10.3	7	41.8	1	7.8	36
South Dakota	17.4	31	29.9	21	30.2	11	54.9	6	7.2	41	36.4	38	7.7	40
Tennessee	22.0	8	32.6	8	32.8	3	64.0	2	10.2	8	32.9	51	8.0	34
Texas	17.9	25	29.1	24	29.5	16	51.8	16	9.3	14	40.8	5	6.4	50
Utah	9.8	51	23.0	49	24.0	42	42.3	48	6.1	49	33.6	50	8.0	34
Vermont	17.1	33	27.0	39	23.3	46	41.9	49	6.2	48	35.7	40	9.9	9
Virginia	19.0	19	27.4	35	25.5	33	48.6	28	8.2	27	36.7	35	7.8	36
Washington	14.9	48	28.0	30	26.9	26	46.3	40	7.7	36	38.7	16	8.8	25
West Virginia	25.5	2	37.6	1	31.7	6	64.8	1	12.3	1	38.4	22	8.8	25
Wisconsin	18.7	20	27.7	32	29.2	17	47.1	35	8.2	27	35.7	40	9.7	14
Wyoming	19.9	17	26.1	44	25.3	37	42.7	46	6.9	44	36.8	34	8.9	24
Range	9.8-25.6		21.5-37.6		18.9-35.3		35.1-60.6		5.7-12.3		32.9-41.8		6.3-11.1	
National Median	17.9		28.6		26.9		49.0		8.3		37.5		8.8	

NOTE: Each percent has a different standard error and confidence interval.
Because of this, rankings are not truly meaningful.
 Unknown and refused responses are *not* included.
 A ranking between 1 and 25 (above the median value) implies poor status relative to other states.
 Data obtained from CDC BRFSS website on 5/18/2010.