

State Prevalence Rankings: BRFSS, 2010

State:	Current Smokers		No Sigmoidoscopy/ Colonoscopy*		Obese Based on BMI		Females Ages 40+ wo/ Mammogram in Past 2 Yrs.		Diabetes		Females 18+ wo/Pap in Past 3 Years		Current Asthma	
	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank
Alabama	21.9	8	36.1	21	33.0	2	24.8	24	13.2	1	16.8	35	8.0	37
Alaska	20.4	13	38.1	17	25.2	38	27.6	19	5.3	51	18.9	26	10.0	7
Arizona	13.5	48	33.2	29	24.7	40	24.4	26	11.4	4	20.7	10	10.0	7
Arkansas	22.9	4	39.1	8	30.9	12	31.3	8	9.6	19	25.2	2	7.8	38
California	12.1	50	38.5	12	24.7	40	21.6	41	8.6	28	19.2	23	7.7	42
Colorado	16.0	33	34.7	26	21.4	51	29.7	11	6.0	50	20.2	13	9.2	22
Connecticut	13.2	49	24.3	51	23.0	48	18.6	47	7.3	41	14.4	46	9.2	22
Delaware	17.3	26	26.0	46	28.7	19	18.6	47	8.7	24	17.3	31	10.0	7
District of Co.	14.8	44	29.2	40	22.7	50	18.2	50	10.9	6	12.9	49	9.9	11
Florida	17.1	28	31.8	34	27.2	28	22.9	33	10.4	8	19.6	19	8.3	34
Georgia	17.6	24	32.3	32	30.4	13	22.8	35	9.7	17	13.4	47	7.8	38
Hawaii	14.5	45	38.5	12	23.1	46	23.5	31	8.3	32	20.3	12	9.4	21
Idaho	15.7	35	40.2	5	26.9	29	36.2	1	7.9	33	23.8	4	8.8	28
Illinois	16.9	30	38.1	17	28.7	19	28.6	15	8.7	24	16.8	35	9.2	22
Indiana	21.2	10	37.2	19	30.2	14	28.7	13	9.8	15	19.8	15	9.5	18
Iowa	16.1	32	35.8	23	29.1	18	24.0	28	7.5	38	19.4	21	7.8	38
Kansas	17.0	29	35.8	23	30.1	15	24.0	28	8.4	31	17.3	31	8.6	31
Kentucky	24.8	2	36.3	20	31.8	5	30.1	9	10.0	14	19.1	24	10.4	4
Louisiana	22.1	7	39.2	7	31.7	6	23.7	30	10.3	10	16.9	33	6.7	50
Maine	18.2	23	25.7	47	27.4	27	19.4	44	8.7	24	15.0	44	10.0	7
Maryland	15.2	40	27.7	44	27.9	22	19.2	46	9.3	21	13.0	48	8.4	32
Massachusetts	14.1	47	24.8	49	23.6	44	16.4	51	7.4	39	11.1	51	10.4	4
Michigan	18.9	18	29.1	41	31.7	6	21.8	40	10.1	12	17.6	30	10.5	3
Minnesota	14.9	43	27.8	43	25.4	37	19.3	45	6.7	48	12.5	50	7.6	43
Mississippi	22.9	4	40.5	4	34.5	1	31.9	7	12.4	2	19.8	15	7.2	49
Missouri	21.1	11	34.8	25	31.4	10	28.7	13	9.4	20	19.9	14	8.8	28
Montana	18.8	19	39.0	9	23.5	45	32.6	5	7.0	45	21.7	6	9.1	26
Nebraska	17.2	27	38.2	16	27.5	26	28.5	17	7.7	36	19.8	15	7.8	38
Nevada	21.3	9	38.5	12	23.1	46	32.8	3	8.5	29	21.6	7	9.2	22
New Hampshire	16.9	30	24.8	49	25.5	36	19.6	43	7.9	33	16.4	38	10.4	4
New Jersey	14.4	46	34.4	27	24.8	39	22.7	36	9.2	22	15.9	42	8.7	30
New Mexico	18.5	20	38.7	11	25.6	35	29.0	12	8.5	29	19.6	19	9.7	15
New York	15.5	37	28.8	42	24.5	42	22.4	37	8.9	23	16.4	38	9.8	13
North Carolina	19.8	15	30.4	37	28.6	21	22.9	33	9.8	15	16.0	41	7.5	44
North Dakota	17.4	25	38.9	10	27.9	22	24.8	24	7.4	39	19.7	18	7.4	46
Ohio	22.5	6	36.0	22	29.7	16	25.8	21	10.1	12	18.3	28	9.6	16
Oklahoma	23.7	3	42.0	2	31.3	11	32.5	6	10.4	8	22.5	5	9.5	18
Oregon	15.1	42	33.9	28	27.6	25	28.6	15	7.2	42	25.1	3	9.5	18
Pennsylvania	18.4	22	32.0	33	29.2	17	26.1	20	10.3	10	18.7	27	9.9	11
Rhode Island	15.7	35	25.5	48	26.0	33	18.6	47	7.8	35	16.9	33	10.9	2
South Carolina	21.0	12	32.7	31	32.0	4	25.5	22	10.7	7	16.1	40	8.3	34
South Dakota	15.4	38	32.9	30	27.7	24	23.4	32	6.9	46	19.1	24	7.5	44
Tennessee	20.1	14	39.5	6	31.7	6	24.3	27	11.3	5	16.6	37	6.0	51
Texas	15.8	34	38.4	15	31.7	6	29.9	10	9.7	17	20.6	11	7.4	46
Utah	9.1	51	30.1	38	23.0	48	33.0	2	6.5	49	26.8	1	9.1	26
Vermont	15.4	38	26.9	45	23.9	43	21.9	39	6.8	47	18.1	29	11.1	1
Virginia	18.5	20	30.7	35	26.4	31	22.2	38	8.7	24	14.8	45	8.4	32
Washington	15.2	40	29.3	39	26.2	32	25.4	23	7.6	37	19.3	22	9.6	16
West Virginia	26.8	1	45.6	1	32.9	3	27.7	18	11.7	3	21.0	9	7.3	48
Wisconsin	19.1	17	30.6	36	26.9	29	21.3	42	7.1	44	15.2	43	8.3	34
Wyoming	19.5	16	40.7	3	25.7	34	32.7	4	7.2	42	21.6	7	9.8	13
Range	9.1-26.8		24.3-45.6		21.4-34.5		16.4-36.2		5.3-13.2		19.0		6.0-11.1	
National Median	17.3		34.8		27.6		24.4		8.7		11.1-26.8		9.1	

*Adults ages 50+ years

NOTE: Each percent has a different standard error and confidence interval.

Because of this, rankings are not truly meaningful.

Unknown and refused responses are not included.

A ranking between 1 and 25 (above the median value) implies poor status relative to other states.

Data obtained from CDC BRFSS prevalence website on 6/7/2011.