

State Prevalence Rankings*: BRFSS, 2011

State	Current Smokers		No Health Care Coverage		Obese Based on BMI		Binge Drinking		Diabetes		<150 Minutes Phys. Activity/Week		High Blood Pressure	
	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank
Alabama	24.3	10	18.3	23	32.0	4	13.7	48	11.8	4	57.6	4	40.1	1
Alaska	22.9	16	20.3	14	27.4	28	20.8	10	7.9	47	42.1	46	29.4	37
Arizona	19.2	37	18.5	21	24.7	40	17.6	33	9.6	24	47.7	28	28.1	47
Arkansas	27.0	3	23.8	5	30.9	7	14.1	47	11.2	6	54.3	7	35.8	8
California	13.7	50	18.4	22	23.8	46	18.6	23	8.9	34	41.8	47	27.8	48
Colorado	18.3	42	17.9	26	20.7	51	20.1	14	6.7	50	38.2	51	25.0	50
Connecticut	17.1	47	12.5	44	24.5	42	17.9	29	9.3	31	47.4	30	29.8	36
Delaware	21.8	23	10.7	47	28.8	19	20.3	13	9.7	22	51.5	16	34.8	10
District of Columbia	20.8	30	7.7	50	23.8	46	25.0	1	9.1	33	42.4	45	30.0	32
Florida	19.3	36	23.6	6	26.6	32	17.1	35	10.4	12	47.2	32	34.2	12
Georgia	21.2	26	25.0	4	28.0	24	16.6	37	10.2	15	49.3	23	32.4	17
Hawaii	16.8	48	9.7	48	21.9	50	21.5	9	8.4	38	41.5	48	28.7	43
Idaho	17.2	46	22.1	8	27.1	29	16.6	37	9.4	30	42.8	43	29.4	37
Illinois	20.9	27	17.1	28	27.1	29	23.0	5	9.7	22	48.3	26	31.0	24
Indiana	25.6	7	19.8	16	30.8	8	17.8	31	10.2	15	54.0	8	32.8	15
Iowa	20.4	31	11.6	45	29.0	18	23.1	4	8.2	42	52.4	14	29.9	34
Kansas	22.0	21	16.7	29	29.6	14	17.0	36	9.5	26	53.2	10	30.8	26
Kentucky	29.0	1	18.6	19	30.4	10	16.1	42	10.8	10	53.2	10	38.0	5
Louisiana	25.7	6	22.5	7	33.4	2	16.1	42	11.8	4	58.0	3	38.4	4
Maine	22.8	18	12.6	43	27.8	25	17.3	34	9.6	24	43.3	42	32.2	19
Maryland	19.1	39	13.0	39	28.3	22	18.0	28	9.5	26	51.3	17	31.3	21
Massachusetts	18.2	43	6.7	51	22.7	49	20.6	12	8.0	45	43.7	41	29.2	40
Michigan	23.3	11	15.2	31	31.3	5	19.7	16	10.0	19	46.5	35	34.2	12
Minnesota	19.1	39	11.4	46	25.7	36	22.1	7	7.3	49	46.0	36	26.3	49
Mississippi	26.0	5	25.7	3	34.9	1	14.2	46	12.4	1	60.0	2	39.3	2
Missouri	25.0	9	17.5	27	30.3	12	19.2	19	10.2	15	50.5	21	34.3	11
Montana	22.1	20	19.8	16	24.6	41	20.8	10	8.0	45	44.7	38	30.2	31
Nebraska	20.0	32	15.9	30	28.4	21	22.7	6	8.4	38	51.1	19	28.6	44
Nevada	22.9	16	27.3	2	24.5	42	18.6	23	10.3	14	47.4	30	30.8	26
New Hampshire	19.4	35	13.4	38	26.2	35	18.7	22	8.7	37	43.9	40	30.7	28
New Jersey	16.8	48	15.0	33	23.7	48	18.2	27	8.8	36	46.8	34	30.6	30
New Mexico	21.5	25	21.8	10	26.3	34	16.4	41	10.0	19	47.8	27	28.4	46
New York	18.1	44	14.7	34	24.5	42	19.6	18	10.5	11	48.5	24	30.7	28
North Carolina	21.8	23	20.7	13	29.1	17	15.2	45	10.9	9	53.2	10	32.4	17
North Dakota	21.9	22	13.6	37	27.8	25	23.8	3	8.2	42	52.7	13	29.1	41
Ohio	25.1	8	14.0	36	29.7	13	20.1	14	10.0	19	48.4	25	32.7	16
Oklahoma	26.1	4	21.9	9	31.1	6	16.5	39	11.1	8	55.2	6	35.5	9
Oregon	19.7	34	18.6	19	26.7	31	16.5	39	9.3	31	38.9	50	29.9	34
Pennsylvania	22.4	19	12.9	40	28.6	20	18.3	26	9.5	26	50.6	20	31.4	20
Rhode Island	20.0	32	14.2	35	25.4	37	19.7	16	8.4	38	51.3	17	32.9	14
South Carolina	23.1	12	20.9	12	30.8	8	15.4	44	12.1	2	50.0	22	36.4	7
South Dakota	23.0	13	12.8	41	28.1	23	22.1	7	9.5	26	53.9	9	31.0	24
Tennessee	23.0	13	18.0	24	29.2	15	10.0	51	11.2	6	61.1	1	38.7	3
Texas	19.2	37	29.8	1	30.4	10	18.9	20	10.2	15	51.8	15	31.3	21
Utah	11.8	51	19.0	18	24.4	45	12.0	49	6.7	50	44.2	39	22.9	51
Vermont	19.1	39	9.2	49	25.4	37	18.5	25	7.7	48	40.8	49	29.3	39
Virginia	20.9	27	15.1	32	29.2	15	17.9	29	10.4	12	47.6	29	31.2	23
Washington	17.5	45	18.0	24	26.5	33	17.8	31	8.9	34	45.8	37	30.0	32
West Virginia	28.6	2	20.2	15	32.4	3	10.1	50	12.0	3	57.0	5	37.0	6
Wisconsin	20.9	27	12.7	42	27.7	27	24.3	2	8.4	38	42.6	44	28.9	42
Wyoming	23.0	13	21.0	11	25.0	39	18.9	20	8.2	42	46.9	33	28.6	44
Range	11.8-29.0		6.7-29.8		20.7-34.9		10.0-25.0		6.7-12.4		38.2-61.1		22.9-40.1	
National Median	21.2		17.9		27.8		18.3		9.5		48.3		30.8	

*Prevalence weighted to landline and cellphone households.
 NOTE: Each percent has a different standard error and confidence interval.
Because of this, rankings are not truly meaningful
 Unknown and refused responses are not included.
 A ranking between 1 and 25 (above the median value) implies poor status relative to other states.
 Data obtained from CDC BRFSS Prevalence and Trends Data accessed August 29, 2012.